

Quantum correlations from incomplete classical statistics

C. Wetterich¹

Institut für Theoretische Physik
Universität Heidelberg
Philosophenweg 16, D-69120 Heidelberg

February 1, 2008

Abstract

We formulate incomplete classical statistics for situations where the knowledge about the probability distribution outside a local region is limited. The information needed to compute expectation values of local observables can be collected in a quantum mechanical state vector, whereas further statistical information about the probability distribution outside the local region becomes irrelevant. The translation of the available information between neighboring local regions is expressed by a Hamilton operator. A quantum mechanical operator can be associated to each local observable, such that expectation values of “classical” observables can be computed by the usual quantum mechanical rules. The requirement that correlation functions should respect equivalence relations for local observables induces a non-commutative product in classical statistics, in complete correspondence to the quantum mechanical operator product. We also discuss the issue of interference and the complex structure of quantum mechanics within our classical statistical setting.

¹e-mail: C.Wetterich@thphys.uni-heidelberg.de

1 Incomplete classical statistics

Following Einstein, Rosen and Podolski [1], many physicists have asked if it could be possible to derive quantum mechanics from an underlying classical statistical system. The motivation is to find an explanation of the “why” of the basic principles of quantum mechanics like the formulation in terms of states and non-commuting operators, the superposition of “probability amplitudes” and the associated interference effects. Such a derivation from an underlying theory could also open the way to possible generalizations of quantum mechanics and to the formulation of further tests of its basic principles [2], [3]. For a large class of classical statistical systems with certain locality properties it has been shown [4] that classical and quantum statistics lead to a different behavior of measurable correlation functions. The experimental verification of the predictions of quantum mechanics is unequivocal. Any attempt to derive quantum mechanics from classical statistics has therefore to circumvent the assumptions of Bell’s inequalities for the classical system and should reproduce the predictions from quantum mechanical interference. Our formulation of incomplete classical statistics with infinitely many degrees of freedom is of this type.

The understanding of classical statistical systems with infinitely many degrees of freedom has made tremendous progress in the past decades. Important links to quantum mechanics have been established by the use of path integrals [5] for the description of the quantum mechanical evolution. This process has diminished considerably the distance between classical statistics and quantum mechanics or quantum field theory [6]. One wonders if quantum mechanics cannot be understood as a particular structure of classical statistical systems with infinitely many degrees of freedom or, in other words, if it can be derived from general statistics [7]. Within general statistics the notions of distance, geometry and topology have already been formulated in terms of properties of correlation functions [8]. The question arises if the notions of time and quantum mechanical evolution can find their origin within the same framework.

The formulation of the basic partition function for classical statistical systems with infinitely many degrees of freedom uses implicitly an assumption of “completeness of the statistical information”. This means that we assign a probability to everyone of the infinitely many configurations. The specification of the probability distribution contains therefore an “infinite amount of information”. This contrasts with the simple observation that only a finite amount of information is available in practice for the computation of the outcome of any physical measurement. A concentration on measurable quantities suggests that the assumption of completeness of the statistical information may have to be abandoned. In this note we explore consequences of “incomplete statistics” which deals with situations where only partial information about the probability distribution is available. In particular, we consider extended systems for which only local information about the probability distribution is given. We will see that the quantum mechanical concepts of states, operators, evolution and interference emerge naturally in this setting.

The outcome of the present work is still far from reaching a stage where quantum mechanics can be derived from a classical statistical setting. The main obstacle remains the characteristic complex structure of quantum mechanics which is not yet implemented in a satisfactory way. Nevertheless, we find it interesting to observe how other structures which were believed to be characteristic for quantum mechanics, like a non-commutative product or the effect of interference, are actually already present in classical statistics. The formulation of incomplete statistics will be very useful in order to motivate how the quantum mechanical structures arise naturally in classical statistics. The notion of incompleteness is, however, not crucial for the existence of these structures. Quantum mechanical structures are present in “standard” (complete) classical statistics as well.

As an example, let us consider a classical statistical system where the infinitely many degrees of freedom φ_n ($n \in \mathbb{Z}$) are ordered in an infinite chain. We concentrate on a “local region” $|\tilde{n}| < \bar{n}$ and assume that the probability distribution $p[\varphi]$ has a “locality property” in the sense that the relative probabilities for any two configurations of the “local variables” $\varphi_{\tilde{n}}$ are independent of the values that take the “external variables” φ_m with $|m| > \bar{n}$. Furthermore, we assume that the probability distribution for the $\varphi_{\tilde{n}}$ is known for given values of the variables $\varphi_{\bar{n}}, \varphi_{-\bar{n}}$ at the border of the local interval. As an example, we may consider a probability distribution

$$\begin{aligned} p[\varphi] &= p_{>}[\varphi_{m \geq \bar{n}}] p_0[\varphi_{-\bar{n} \leq \tilde{n} \leq \bar{n}}] p_{<}[\varphi_{m \leq -\bar{n}}] \\ p_0[\varphi] &= \exp \left\{ - \sum_{|\tilde{n}| < \bar{n}} \left[\frac{\epsilon}{2} \mu^2 \varphi_{\tilde{n}}^2 + \frac{\epsilon}{8} \lambda \varphi_{\tilde{n}}^4 + \frac{M}{2\epsilon} (\varphi_{\tilde{n}} - \varphi_{\tilde{n}-1})^2 \right] - \frac{M}{2\epsilon} (\varphi_{\bar{n}} - \varphi_{\bar{n}-1})^2 \right\} \end{aligned} \quad (1.1)$$

where $p_{>}$ and $p_{<}$ are only constrained by the overall normalization of $p[\varphi]$ and we will consider the limit $\epsilon \rightarrow 0$. This statistical system cannot be reduced to a system with a finite number of degrees of freedom since the probability for the occurrence of specific values of the “border variables” $\varphi_{\bar{n}}, \varphi_{-\bar{n}}$ depends on the values of the external variables φ_m and their probability distribution. The statistical information about this system is incomplete since we will not specify the probability distribution $p_{>} p_{<}$ for the external variables φ_m completely.

Local observables are constructed from the local variables $\varphi_{\tilde{n}}$. As usual, their expectation values are computed by “functional integrals” where the probability distribution $p[\varphi]$ appears as a weight factor. Incomplete statistics deal with the expectation values of local observables. Thereby the questions that may be asked reflect the incompleteness of the statistical information. As a first question, we ask what are the conditions on $p_{>} p_{<}$ which lead to $\tilde{n}$ -independent expectation values $\langle \varphi^p(\tilde{n}) \rangle$ for all or some integer values of p ? Furthermore, are there any restrictions on the allowed values of $\tilde{n}$ -independent expectation values $\langle \varphi^p(\tilde{n}) \rangle$? The answer to these questions, given in sect. 4, may surprise: indeed, probability distributions $p_{>}, p_{<}$ that are consistent with $\tilde{n}$ -independent $\langle \varphi^p(\tilde{n}) \rangle$ are possible only for certain discrete values for $\langle \varphi^p(\tilde{n}) \rangle$. These values correspond precisely

to the expectation values of the operators $\hat{Q}^p$ in the stationary states of quantum mechanics. Here our example (1.1) corresponds to a one-dimensional particle with mass M in an unharmonic potential $V(\hat{Q}) = \frac{\mu^2}{2}\hat{Q}^2 + \frac{\lambda}{8}\hat{Q}^4$. The appearance of the quantum-mechanical discreteness in a classical statistical system is striking and we will attempt to understand its origin and the deep connection to quantum mechanics in more detail in this note.

We next ask more generally what is the minimal amount of information about the probability distribution for the external variables φ_m which is necessary for a computation of expectation values of local observables. One finds that this information can be summarized in “states” $|\psi\rangle, \langle\psi|$ that can be represented as ordinary functions $\{\psi(\varphi_{\tilde{n}}), \psi(\varphi_{-\tilde{n}})\}$. Since these functions depend each only on one variable, the specification of the states contains much less information than the full probability distribution $p_{>}p_{<}$ which depends on infinitely many variables $\varphi_{m\geq\tilde{n}}, \varphi_{m\leq-\tilde{n}}$. The states contain the minimal information for “local questions” and are therefore the appropriate quantities for our formulation of incomplete statistics. We will see in sect. 7 that any further information about the probability distributions $p_{>}[\varphi_m], p_{<}[\varphi_m]$ beyond the one contained in the state vectors is actually irrelevant for the computation of expectation values of local observables.

The expectation values of all local observables can be computed from the knowledge of the local probability distribution and the states $|\psi\rangle$ and $\langle\psi|$. For this computation one associates to every local observable $A[\varphi]$ an appropriate operator $\hat{A}$ and finds the prescription familiar from quantum mechanics

$$\langle A[\varphi] \rangle = \langle \psi | \hat{A} | \psi \rangle \quad (1.2)$$

There is a unique mapping $A[\varphi] \rightarrow \hat{A}$ for every local observable which can be expressed in terms of an appropriate functional integral. We find that for simple observables $A[\varphi]$ the operators $\hat{A}$ correspond precisely to familiar operators in quantum mechanics. For example, the observable $\varphi(\tilde{n})$ can be associated to the operator $\hat{Q}(\tau)$ in the Heisenberg picture where time is analytically continued, $\tau = it$, and $\tilde{n} = \tau/\epsilon$.

Local correlation functions involving derivatives may be ambiguous in the continuum limit. This problem is well known in functional integral formulations of quantum field theories. We show how to avoid this problem by defining correlations in terms of equivalence classes of observables. In fact, two observables $A_1[\varphi], A_2[\varphi]$ can sometimes be represented by the same operator $\hat{A}$. In this case $A_1[\varphi]$ and $A_2[\varphi]$ are equivalent since they cannot be distinguished by their expectation values for arbitrary states. They have the same expectation values for all possible probability distributions. We define a product between equivalence classes of observables which can be associated to the product of operators. For example, we associate to a correlation $\varphi(\tilde{n}_1) \circ \varphi(\tilde{n}_2)$ the product $\hat{Q}(\tau_1)\hat{Q}(\tau_2)$ which is not commutative. It is again striking how the non-commutativity of quantum mechanics arises directly from the question what are meaningful correlation functions in incomplete statistics! Indeed, equivalent observables should lead to equivalent correlations. We will see (sect. 6)

that this points to a non-commutative definition of the correlation $\varphi(\tilde{n}_1) \circ \varphi(\tilde{n}_2)$. In sect. 7 we argue that the “quantum correlation” based on $\varphi_1 \circ \varphi_2$ can actually be formulated quite generally in classical statistics (not necessarily incomplete) and has better “robustness properties” as compared to the usual classical correlation.

We also may ask what can be learned for expectation values like $\langle \varphi^p(\tilde{n}) \rangle$ if the states are specified by a sufficient number of expectation values at some given location $\tilde{n}_0$ (e.g. $\langle \varphi^p(\tilde{n}_0) \rangle$, $\langle (\varphi(\tilde{n}_0+1) - \varphi(\tilde{n}_0))^{p'} \rangle$ etc.). We find that operators and states can be transported from one site to the next by an evolution operator $\hat{U}$ in complete analogy to quantum mechanics. This introduces once more a crucial concept of quantum mechanics in the formulation of incomplete statistics, namely the Hamilton operator. As one may expect, it corresponds to the transfer matrix in classical statistics.

Finally, we ask in sect. 8 if the characteristic interference of quantum mechanics also finds its equivalent in incomplete statistics. We explicitly construct classical probability distributions that correspond to linear superpositions of states $\alpha_1|\psi_1\rangle + \alpha_2|\psi_2\rangle$ and show the corresponding interference behavior. We also describe probability distributions leading to a formulation in terms of density matrices that do not correspond to pure states. We conclude in sect. 9 that many fundamental features of quantum mechanics are shown to arise directly from the formulation of incomplete statistics. Inverting our procedure we find quantum mechanical states which admit a functional integral description. This goes beyond the usual description of the vacuum or thermal equilibrium state.

There remains one crucial feature of quantum mechanics, namely its complex structure or the issue of “phases”, that has not yet emerged in a satisfactory way from classical statistics. It is closely related to the difference between “real time” in quantum mechanics and “Euclidean time” in classical statistics or the issue of analytical continuation. We describe our (incomplete) attempts in this direction in two appendices. The invariance of the probability distribution under a reflection $\varphi_n \rightarrow \varphi_{-n}$ can be associated with a complex structure. With respect to this complex structure all quantities that are odd under reflections become purely imaginary. Then the evolution operators $\hat{U}$ are unitary and $\langle \psi |$ is complex conjugate to $|\psi\rangle$. Also $\tau = \epsilon n$ is imaginary. It is not clear, however, how the state vectors as functions of φ_n (at fixed n) should be analytically continued without losing the probability interpretation of the functional integral.

2 States and operators

Consider a discrete ordered set of continuous variables $\varphi_n \equiv \varphi(\tau)$, $\tau = \epsilon n$, $n \in \mathbb{Z}$ and a normalized probability distribution $p(\{\varphi_n\}) \equiv p[\varphi] = \exp(-S[\varphi])$ with $\int D\varphi e^{-S[\varphi]} \equiv \prod_n (\int_{-\infty}^{\infty} d\varphi_n) p[\varphi] = 1$. We will assume that the action S is local in a

range $-\bar{\tau} < \tau < \bar{\tau}$, i.e.

$$\begin{aligned} S &= -\ln p = \int_{-\bar{\tau}}^{\bar{\tau}} d\tau' \mathcal{L}(\tau') + S_{>}(\bar{\tau}) + S_{<}(-\bar{\tau}) \\ \mathcal{L}(\tau') &= V(\varphi(\tau'), \tau') + \frac{1}{2} Z(\tau') (\partial_{\tau'} \varphi(\tau'))^2 \end{aligned} \quad (2.1)$$

Here we have used a continuum notation ($n_{1,2} = \tau_{1,2}/\epsilon$) which can be translated into a discrete language by

$$\begin{aligned} \int_{\tau_1}^{\tau_2} d\tau' \mathcal{L}(\tau') &= \epsilon \sum_{n=n_1+1}^{n_2-1} \mathcal{L}_n + \frac{\epsilon}{2} [V_{n_2}(\varphi_{n_2}) + V_{n_1}(\varphi_{n_1})] \\ &+ \frac{\epsilon}{4} \left[Z_{n_2} \left(\frac{\varphi_{n_2} - \varphi_{n_2-1}}{\epsilon} \right)^2 + Z_{n_1} \left(\frac{\varphi_{n_1+1} - \varphi_{n_1}}{\epsilon} \right)^2 \right] \end{aligned} \quad (2.2)$$

with

$$\mathcal{L}_n = V_n(\varphi_n) + \frac{Z_n}{4\epsilon^2} \{(\varphi_{n+1} - \varphi_n)^2 + (\varphi_n - \varphi_{n-1})^2\} \quad (2.3)$$

This corresponds to a discrete derivative

$$\begin{aligned} (\partial_{\tau} \varphi(\tau))^2 &= \frac{1}{2} \left\{ \left(\frac{\varphi(\tau + \epsilon) - \varphi(\tau)}{\epsilon} \right)^2 + \left(\frac{\varphi(\tau) - \varphi(\tau - \epsilon)}{\epsilon} \right)^2 \right\} \\ &= \frac{1}{2\epsilon^2} \{(\varphi_{n+1} - \varphi_n)^2 + (\varphi_n - \varphi_{n-1})^2\}. \end{aligned} \quad (2.4)$$

The boundary terms in eq. (2.2) are chosen such that $S_{>}(\bar{\tau})$ is independent of all $\varphi(\tau')$ with $\tau' < \bar{\tau}$ whereas $S_{<}(-\bar{\tau})$ only depends on $\varphi(\tau' \leq -\bar{\tau})$. Except for the overall normalization of p no additional assumptions about the form of $S_{>}(\bar{\tau})$ and $S_{<}(-\bar{\tau})$ will be made. In case of S being local also at $\bar{\tau}$ we note that $S_{>}(\bar{\tau})$ contains a term $\frac{\epsilon}{2} [V(\varphi(\bar{\tau}), \bar{\tau}) + V(\varphi(\bar{\tau} + \epsilon), \bar{\tau} + \epsilon)] + \frac{\epsilon}{4} (Z(\bar{\tau}) + Z(\bar{\tau} + \epsilon)) \left(\frac{\varphi(\bar{\tau} + \epsilon) - \varphi(\bar{\tau})}{\epsilon} \right)^2$, which involves a product $\varphi(\bar{\tau} + \epsilon)\varphi(\bar{\tau})$ and therefore links the variables with $\tau > \bar{\tau}$ to the ones with $\tau \leq \bar{\tau}$.

We are interested in local observables $A[\varphi; \tau]$ which depend only on those $\varphi(\tau')$ where $\tau - \frac{\delta}{2} \leq \tau' \leq \tau + \frac{\delta}{2}$. (We assume $-\bar{\tau} < \tau - \frac{\delta}{2}$, $\bar{\tau} > \tau + \frac{\delta}{2}$.) As usual, the expectation value of A is

$$\langle A(\tau) \rangle = \int D\varphi A[\varphi; \tau] e^{-S[\varphi]} \quad (2.5)$$

As mentioned in the introduction, our investigation concerns the question what we can learn about expectation values of local observables and suitable products thereof in a situation where we have no or only partial information about $S_{>}(\bar{\tau})$ and $S_{<}(-\bar{\tau})$. It seems obvious that the full information contained in S is not needed if only

expectation values of local observables of the type (2.5) are to be computed. On the other hand, $\langle A(\tau) \rangle$ cannot be completely independent of $S_>(\bar{\tau})$ and $S_<(-\bar{\tau})$ since the next neighbor interactions (2.2) relate “local variables” $\varphi(\tau - \frac{\delta}{2} < \tau' < \tau + \frac{\delta}{2})$ to the “exterior variables” $\varphi(\tau' > \bar{\tau})$ and $\varphi(\tau' < -\bar{\tau})$.

In order to establish the necessary amount of information needed from $S_>(\bar{\tau})$ and $S_<(-\bar{\tau})$ we first extend $S_>$ and $S_<$ to values $|\tau| < \bar{\tau}$

$$S_<(\tau_1) = S_<(-\bar{\tau}) + \int_{-\bar{\tau}}^{\tau_1} d\tau' \mathcal{L}(\tau'), \quad S_>(\tau_2) = S_>(\bar{\tau}) + \int_{\tau_2}^{\bar{\tau}} d\tau' \mathcal{L}(\tau') \quad (2.6)$$

where we note the general identity

$$S_>(\tau) + S_<(\tau) = S \quad (2.7)$$

The expectation value (2.5) can be written as

$$\begin{aligned} \langle A(\tau) \rangle &= \int d\varphi(\tau + \frac{\delta}{2}) \int d\varphi(\tau - \frac{\delta}{2}) \int D\varphi_{(\tau' > \tau + \frac{\delta}{2})} e^{-S_>(\tau + \frac{\delta}{2})} \\ &\int D\varphi_{(\tau - \frac{\delta}{2} < \tau' < \tau + \frac{\delta}{2})} A[\varphi; \tau] \exp\left\{-\int_{\tau - \frac{\delta}{2}}^{\tau + \frac{\delta}{2}} d\tau'' \mathcal{L}(\tau'')\right\} \int D\varphi_{(\tau' < \tau - \frac{\delta}{2})} e^{-S_<(\tau - \frac{\delta}{2})} \end{aligned} \quad (2.8)$$

This suggests the introduction of the “states”

$$\begin{aligned} |\psi(\varphi(\tau - \frac{\delta}{2}); \tau - \frac{\delta}{2})\rangle &= \int D\varphi_{(\tau' < \tau - \frac{\delta}{2})} e^{-S_<(\tau - \frac{\delta}{2})} \\ \langle \psi(\varphi(\tau + \frac{\delta}{2}); \tau + \frac{\delta}{2})| &= \int D\varphi_{(\tau' > \tau + \frac{\delta}{2})} e^{-S_>(\tau + \frac{\delta}{2})} \end{aligned} \quad (2.9)$$

and the operator

$$\begin{aligned} \hat{A}_\delta(\varphi(\tau + \frac{\delta}{2}), \varphi(\tau - \frac{\delta}{2}); \tau) \\ = \int D\varphi_{(\tau - \frac{\delta}{2} < \tau' < \tau + \frac{\delta}{2})} A[\varphi; \tau] \exp\left\{-\int_{\tau - \frac{\delta}{2}}^{\tau + \frac{\delta}{2}} d\tau'' \mathcal{L}(\tau'')\right\} \end{aligned} \quad (2.10)$$

We note that $|\psi\rangle$ is a function of $\varphi(\tau - \frac{\delta}{2})$ since the latter appears in $S_<(\tau - \frac{\delta}{2})$ and is not included in the (“functional”) integration (2.9). Similarly, $\langle \psi|$ depends on $\varphi(\tau + \frac{\delta}{2})$ whereas $\hat{A}$ is a function of the two variables $\varphi(\tau + \frac{\delta}{2})$ and $\varphi(\tau - \frac{\delta}{2})$. Using a notation where $|\psi\rangle$ and $\langle \psi|$ are interpreted as (infinite dimensional) vectors and $\hat{A}$ as a matrix, one has

$$\begin{aligned} \langle A(\tau) \rangle &= \langle \psi(\tau + \frac{\delta}{2}) | \hat{A}_\delta(\tau) | \psi(\tau - \frac{\delta}{2}) \rangle \\ &\equiv \int d\varphi_2 \int d\varphi_1 \langle \psi(\varphi_2; \tau + \frac{\delta}{2}) | \hat{A}_\delta(\varphi_2, \varphi_1; \tau) | \psi(\varphi_1; \tau - \frac{\delta}{2}) \rangle \end{aligned} \quad (2.11)$$

This form resembles already the well-known prescription for expectation values of operators in quantum mechanics. In contrast to quantum mechanics eq. (2.11) still involves, however, two different state vectors.

The mapping $A[\varphi; \tau] \rightarrow \hat{A}_\delta(\tau)$ can be computed (cf. eq. (2.10)) if $\mathcal{L}(\tau')$ is known for $|\tau'| < \bar{\tau}$. The only information needed from $S_>(\bar{\tau})$ and $S_<(-\bar{\tau})$ is therefore contained in the two functions $\{\psi(\varphi)|$ and $|\psi(\varphi)\}$! The specification of these states (wave functions) at $\bar{\tau}$ and $-\bar{\tau}$ and of $\mathcal{L}(|\tau| < \bar{\tau})$ completely determines the expectation values of *all* local observables!

We will see below the close connection to the states in quantum mechanics. In our context we emphasize that for any given S these states can be computed as well defined functional integrals (2.9). Due to eq. (2.7) they obey the normalization

$$\{\psi(\tau)\psi(\tau)\} \equiv \int d\varphi \{\psi(\varphi; \tau)|\psi(\varphi; \tau)\} = 1 \quad (2.12)$$

Incomplete statistics explores statements that can be made for local observables and appropriate products thereof without using information about $S_>$ or $S_<$ beyond the one contained in the states $|\psi\rangle$ and $\langle\psi|$.

3 Evolution in Euclidean time

For a “locality interval” $\delta > 0$ the expression (2.11)) involves states at different locations or “Euclidean times” $\tau + \frac{\delta}{2}$ and $\tau - \frac{\delta}{2}$. We aim for a formulation where only states at the same τ appear. We therefore need the explicit mapping from $|\psi(\tau - \frac{\delta}{2})\}$ to a reference point $|\psi(\tau)\}$ and similar for $\{\psi(\tau + \frac{\delta}{2})|$. This mapping should also map $\hat{A}_\delta$ to a suitable operator such that the structure (2.11) remains preserved. The dependence of states and operators on the Euclidean time τ is described by evolution operators ($\tau_2 > \tau_1$, $\tau_2 > \tau_f$, $\tau_i > \tau_1, \tau_f = \tau + \frac{\delta}{2}$, $\tau_i = \tau - \frac{\delta}{2}$)

$$\begin{aligned} |\psi(\tau_2)\} &= \hat{U}(\tau_2, \tau_1)|\psi(\tau_1)\} \\ \{\psi(\tau_1)| &= \{\psi(\tau_2)|\hat{U}(\tau_2, \tau_1) \\ \hat{A}(\tau_2, \tau_1) &= \hat{U}(\tau_2, \tau_f)\hat{A}(\tau_f, \tau_i)\hat{U}(\tau_i, \tau_1) \end{aligned} \quad (3.1)$$

or differential operator equations ($\epsilon \rightarrow 0$);

$$\begin{aligned} \partial_\tau |\psi(\tau)\} &= -\hat{H}(\tau)|\psi(\tau)\} \\ \partial_\tau \{\psi(\tau)| &= \{\psi(\tau)|\hat{H}(\tau) \\ \partial_{\tau_i} \hat{A}_\delta(\tau_f, \tau_i) &= \hat{A}_\delta(\tau_f, \tau_i)\hat{H}(\tau_i) \\ \partial_{\tau_f} \hat{A}_\delta(\tau_f, \tau_i) &= -\hat{H}(\tau_f)\hat{A}_\delta(\tau_f, \tau_i). \end{aligned} \quad (3.2)$$

The evolution operator has an explicit representation as a functional integral

$$\hat{U}(\varphi(\tau_2), \varphi(\tau_1); \tau_2, \tau_1) = \int D\varphi_{(\tau_1 < \tau' < \tau_2)} \exp \left\{ - \int_{\tau_1}^{\tau_2} d\tau'' \mathcal{L}(\tau'') \right\} \quad (3.3)$$

and obeys the composition property ($\tau_3 > \tau_2 > \tau_1$)

$$\hat{U}(\tau_3, \tau_2)\hat{U}(\tau_2, \tau_1) = \hat{U}(\tau_3, \tau_1) \quad (3.4)$$

with

$$\hat{U}(\varphi_2, \varphi_1; \tau, \tau) = \delta(\varphi_2 - \varphi_1) \quad (3.5)$$

It can therefore be composed as a product of transfer matrices or “infinitesimal” evolution operators

$$\hat{U}(\tau + \epsilon, \tau) = e^{-\epsilon \hat{H}(\tau + \frac{\epsilon}{2})} \quad (3.6)$$

In case of translation symmetry for the local part of the probability distribution, i. e. for V and Z independent of τ , we note the symmetry in $\varphi_1 \leftrightarrow \varphi_2$

$$\hat{U}(\tau + \epsilon, \tau) = \hat{U}^T(\tau + \epsilon, \tau), \quad \hat{H}(\tau + \frac{\epsilon}{2}) = \hat{H}^T(\tau + \frac{\epsilon}{2}) = \hat{H} \quad (3.7)$$

In this case the real symmetric matrix $\hat{H}$ has real eigenvalues E_n . Then the general solution of the differential equation (3.2) may be written in the form

$$|\psi(\tau)\rangle = \sum_n \psi_0^{(n)} e^{-E_n \tau}, \quad \langle \psi(\tau) | = \sum_n \bar{\psi}_0^{(n)} e^{E_n \tau} \quad (3.8)$$

where $\psi_0^{(n)}$ and $\bar{\psi}_0^{(n)}$ are eigenvectors of $\hat{H}$ with eigenvalues E_n . Here we recall that the construction (2.9) implies that $|\psi\rangle$ and $\langle \psi|$ are real positive functions of φ for every τ . This restricts the allowed values of the coefficients $\psi_0^{(n)}, \bar{\psi}_0^{(n)}$.

We next want to compute the explicit form of the Hamilton operator $\hat{H}$. It is fixed uniquely by the functional integral representation (3.3) for $\hat{U}$. In order to obey the defining equation (3.6), the Hamilton operator $\hat{H}$ must fulfill for arbitrary $|\psi(\varphi)\rangle$ the relation (with $Z = Z(\tau + \frac{\epsilon}{2}) = \frac{1}{2}(Z(\tau + \epsilon) + Z(\tau))$)

$$\begin{aligned} \int d\varphi_1 \hat{H}(\varphi_2, \varphi_1) |\psi(\varphi_1)\rangle &= -\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \left\{ \int d\varphi_1 \right. \\ &\quad \left. \exp \left[-\frac{\epsilon}{2}(V(\varphi_2) + V(\varphi_1)) - \frac{Z}{2\epsilon}(\varphi_2 - \varphi_1)^2 \right] |\psi(\varphi_1)\rangle - |\psi(\varphi_2)\rangle \right\} \end{aligned} \quad (3.9)$$

The solution of this equation can be expressed in terms of the operators

$$\begin{aligned} \hat{Q}(\varphi_2, \varphi_1) &= \varphi_1 \delta(\varphi_2 - \varphi_1) \\ \hat{P}^2(\varphi_2, \varphi_1) &= -\delta(\varphi_2 - \varphi_1) \frac{\partial^2}{\partial \varphi_1^2} \end{aligned} \quad (3.10)$$

as

$$\hat{H}(\tau) = V(\hat{Q}, \tau) + \frac{1}{2Z(\tau)} \hat{P}^2 \quad (3.11)$$

This can be established² by using under the φ_1 -integral the replacement

$$e^{-\frac{Z}{2\epsilon}(\varphi_2 - \varphi_1)^2} \rightarrow \left(\frac{2\pi\epsilon}{Z}\right)^{1/2} \delta(\varphi_2 - \varphi_1) \exp\left(\frac{\epsilon}{2Z} \frac{\partial^2}{\partial \varphi_1^2}\right) \quad (3.12)$$

which is valid by partial integration if the integrand decays fast enough for $|\varphi_1| \rightarrow \infty$. We note that the operators $\hat{Q}$ and $\hat{P}^2$ do not commute, e.g.

$$[\hat{P}^2, \hat{Q}](\varphi_2, \varphi_1) = -2\delta(\varphi_2 - \varphi_1) \frac{\partial}{\partial \varphi_1} \quad (3.13)$$

The Hamilton operator can be used in order to establish the existence of the inverse of the “infinitesimal” evolution operator, $\hat{U}^{-1}(\tau + \epsilon, \tau) = e^{\epsilon \hat{H}(\tau + \frac{\epsilon}{2})}$. Then the inverse $\hat{U}^{-1}(\tau_2, \tau_1)$ is defined by the multiplication of “infinitesimal” inverse evolution operators, and we can extend the composition property (3.4) to arbitrary τ be defining for $\tau_2 < \tau_1$

$$\hat{U}(\tau_2, \tau_1) = \hat{U}^{-1}(\tau_1, \tau_2) \quad (3.14)$$

(For a given dependence of $\hat{U}$ on the variables τ_2 and τ_1 the matrix $\hat{U}(\tau_1, \tau_2)$ obtains from $\hat{U}(\tau_2, \tau_1)$ by a simple exchange of the arguments τ_1 and τ_2 .) Using eq. (3.1), this allows us to write the expectation value of a local observable in a form involving states at the same τ -variable

$$\langle A(\tau) \rangle = \{\psi(\tau) \hat{U}(\tau, \tau + \frac{\delta}{2}) \hat{A}_\delta(\tau) \hat{U}(\tau - \frac{\delta}{2}, \tau) \psi(\tau)\} \quad (3.15)$$

4 Stationary observables

At this point we can already answer the question posed in the introduction: What are the conditions for the expectation values $\varphi^p(\tau)$ to be τ -independent for all p or some given p ? Since the observables $\varphi(\tau)$ have support only at one point τ , we can choose $\delta = 0$ such that

$$\langle \varphi^p(\tau) \rangle = \{\psi(\tau) \hat{Q}^p \psi(\tau)\} = \int d\varphi \{\psi(\varphi; \tau) |\varphi^p| \psi(\varphi; \tau)\} \quad (4.1)$$

Using the explicit solution of the evolution equation (3.8), this yields for translation symmetric V and Z

$$\langle \varphi^p(\tau) \rangle = \int d\varphi \sum_{n, n'} \bar{a}_n \chi_n(\varphi) e^{E_n \tau} \varphi^p e^{-E_{n'} \tau} a_{n'} \chi_{n'}(\varphi) \quad (4.2)$$

²The Hamilton operator can also be defined for small nonzero ϵ where one has $e^{-\epsilon \hat{H}} = \left(\frac{2\pi\epsilon}{Z}\right)^{1/2} e^{-\frac{\epsilon}{2} V(\hat{Q}, \tau + \epsilon)} e^{-\epsilon \hat{P}^2 / 2Z} e^{-\frac{\epsilon}{2} V(\hat{Q}, \tau)}$. We have absorbed the prefactor by a constant shift in V , i.e. $V \rightarrow V - \frac{1}{2\epsilon} \ln \frac{2\pi\epsilon}{Z}$. If appropriate, one may use the symmetric form where $\partial_1^2 \rightarrow \frac{1}{4}(\vec{\partial}_1 - \vec{\partial}_2)^2$, $\partial_i = \partial / \partial \varphi_i$.

where the functions $\chi_n(\varphi)$ are an orthonormal set of basis vectors. We choose them to be eigenfunctions of the Hamiltonian

$$\hat{H}\chi_n(\varphi) = E_n\chi_n(\varphi) \quad (4.3)$$

where $\hat{H}$ is interpreted as a differential operator which takes for $Z = M$ the familiar form

$$\hat{H} = V(\varphi) - \frac{1}{2M} \frac{\partial^2}{\partial \varphi^2} \quad (4.4)$$

We specialize to $V(\varphi) = \frac{1}{2}\mu^2\varphi^2 + \frac{1}{8}\lambda\varphi^4$, $\lambda > 0$, which corresponds to an unharmonic oscillator (cf. eq. (1.1)). The functions $\chi_n(\varphi)$ are, of course, precisely the stationary solutions of the Schrödinger equation for a particle in a one-dimensional potential $V(\varphi)$.

Defining

$$K_{nn'}^p = \int d\varphi \chi_n(\varphi) \varphi^p \chi_{n'}(\varphi) \quad (4.5)$$

one obtains

$$\langle \varphi^p(\tau) \rangle = \sum_{n,n'} \bar{a}_n a_{n'} K_{nn'}^p e^{(E_n - E_{n'})\tau} \quad (4.6)$$

An obvious solution for τ -independent expectation values $\langle \varphi^p(\tau) \rangle$ are probability distributions for which $a_n, a_{n'}$ are nonvanishing only for one given value $n = n'$. Such distributions correspond precisely to the stationary pure states in the associated quantum mechanical problem. Furthermore, the normalization (2.12) implies $\bar{a}_n a_n = 1$ and therefore

$$\langle \varphi^p(\tau) \rangle = K_{nn}^p \quad (4.7)$$

If the above solution is the only solution, we arrive at a simple, perhaps surprising conclusion: The mere requirement of τ -independent $\langle \varphi^p(\tau) \rangle$ associates the allowed probability distributions to stationary quantum states. Furthermore, the allowed values of $\langle \varphi^p \rangle$ are discrete, reflecting the discreteness of the associated quantum mechanical problem!

It remains to be shown that no other solutions exist with $\partial_\tau \langle \varphi^p(\tau) \rangle = 0$. If we require the τ -independence for $\langle \varphi^p(\tau) \rangle$ for all p , this can be established in a rather straightforward way. Assume $\bar{a}_n \neq 0$ for some n . If a_m is nonvanishing for $m \neq n$, the sum (4.6) contains a τ -dependent contribution $a_n a_m K_{nm}^p \exp\{(E_n - E_m)\tau\}$ at least for certain values of p . (There is no superselection rule for the unharmonic oscillator. Therefore for any given pair (n, m) the integral K_{nm}^p does not vanish for all p .) The above contribution to $\partial_\tau \langle \varphi^p(\tau) \rangle$ could only be cancelled by a similar contribution with the same τ -dependence, whereby $\bar{a}_{n'}$ and $a_{m'}$ must be nonzero and $E_{n'} - E_{m'} = E_n - E_m$, with $n' \neq n$, $m' \neq m$. The spectrum of the unharmonic oscillator has no degeneracy – there is only one state for every eigenvalue E_n . The equality of two energy differences in the spectrum could occur accidentally for particular values of λ . This is, however, not a problem for our argument since a cancellation of the term $\sim \exp\{(E_n - E_m)\tau\}$ would only be possible

at the expense of new terms with a nontrivial τ -dependence, i.e. $\sim \exp\{(E_n - E_{m'})\tau\}$ and $\exp(E_{n'} - E_m)\tau$. This problem will be continued if one tries further to cancel the new terms. Since equal energy differences can only occur accidentally, one rapidly arrives at a contradiction since no pair with the required precise energy difference will be available any more. (A similar argument would not work so easily for the harmonic oscillator with its equidistant spectrum.)

Finally, if we require τ -dependence only for some values of p , the above argument will work only if for every pair (m, n) there exists a nonvanishing K_{mn}^p . Otherwise it is easy to construct a counterexample. Assume we consider only $p = 2$ and there is, perhaps accidentally, a pair (m, n) with $K_{mn}^2 = 0$. Then a state $|\psi\rangle \sim (a_m\chi_m + a_n\chi_n)$ and similar for $\langle\psi|$ leads also to τ -independent values of $\langle\varphi^2(\tau)\rangle$. The discreteness of the allowed values of τ -independent $\langle\varphi^2\rangle$ would be lost in this case. However, if a sufficient number of expectation values $\langle\varphi^p(\tau)\rangle$ are required to be τ -independent simultaneously, this possibility of accidentally vanishing matrix elements can be discarded.

To conclude, the condition of τ -independence of a suitable number of expectation values $\langle\varphi^p(\tau)\rangle$ leaves only a discrete number of possibilities to continue the local probability distribution (1.1) outside the local region. Correspondingly, the allowed τ -independent values $\langle\varphi^p\rangle$ are also discrete. They correspond to the stationary wave functions of the quantum mechanical unharmonic oscillator. Since, in addition, $|\psi(\varphi)\rangle$ is real and positive one concludes that excited states with zeros of $|\psi(\varphi)\rangle$ cannot be described in our setting. The ground state solution can be realized by extending the translation symmetry of p_0 also to $p_>$ and $p_<$, but also by all other $p_>$ and $p_<$ which lead to the same state vectors $|\psi\rangle$ and $\langle\psi|$. We have seen that the simple question of τ -independence of expectation values has led us directly to the appearance of quantum mechanical concepts like states and operators in our classical statistical setting. In the following we will exploit these quantum mechanical concepts to describe also τ -dependent expectation values. This will give us a deeper insight how various quantum mechanical structures are rooted in our formulation of incomplete classical statistics.

5 Schrödinger and Heisenberg operators

In this section we want to exploit further the mapping between incomplete statistics and quantum mechanics for situations where expectation values like $\langle\varphi(\tau)\rangle$ may depend on τ . A typical question one may ask within incomplete classical statistics is the following: Given a large set of measurements of observables with support at a given value $\tau = 0$, like $\langle\varphi^p(0)\rangle$, $\langle(\partial_\tau\varphi(0))^{p'}\rangle$, etc., what can one predict for the expectation values of similar observables at some other location $\tau \neq 0$? It is obvious that the evolution operator $\hat{U}$ is the appropriate tool to tackle this type of questions.

The existence of the inverse evolution operator allows us to associate to an observable $A(\tau)$ the operator $\hat{A}_S(\tau)$ in the Schrödinger representation (cf. eq. (3.15))

$$\hat{A}_S(\tau) = \hat{U}(\tau, \tau + \frac{\delta}{2}) \hat{A}_\delta(\tau) \hat{U}(\tau - \frac{\delta}{2}, \tau) \quad (5.1)$$

The expectation value of the observable A can be expressed by the expectation value of the operator $\hat{A}_S$ in a way analogous to quantum mechanics

$$\langle A(\tau) \rangle = \{\psi(\tau) | \hat{A}_S(\tau) | \psi(\tau)\} = \text{Tr} \rho(\tau) \hat{A}_S(\tau) \quad (5.2)$$

For the second identity we have introduced the “density matrix”

$$\begin{aligned} \rho(\varphi_1, \varphi_2, \tau) &= |\psi(\varphi_1, \tau)\{\psi(\varphi_2, \tau)| = \int D\varphi_{(\tau' \neq \tau)} e^{-S(\varphi_1, \varphi_2)} \\ \text{Tr} \rho(\tau) &= 1 \end{aligned} \quad (5.3)$$

where $S(\varphi_1, \varphi_2)$ obtains from S by replacing $\varphi(\tau) \rightarrow \varphi_1$ for all “kinetic” terms involving $\varphi(\tau' < \tau)$ and $\varphi(\tau) \rightarrow \varphi_2$ for those involving $\varphi(\tau' > \tau)$, whereas for potential terms $e^{-\epsilon V(\varphi(\tau))} \rightarrow e^{-\frac{\epsilon}{2}(V(\varphi_1) + V(\varphi_2))}$.

In order to make the transition to the Heisenberg picture, we may select a reference point $\tau = 0$ and define

$$\hat{U}(\tau) \equiv \hat{U}(\tau, 0), \quad \rho \equiv \rho(\tau = 0), \quad \rho(\tau) = \hat{U}(\tau) \rho \hat{U}^{-1}(\tau) \quad (5.4)$$

This specifies the Heisenberg picture for the τ -dependent operators

$$\begin{aligned} \hat{A}_H(\tau) &= \hat{U}^{-1}(\tau) \hat{A}_S(\tau) \hat{U}(\tau) \\ \langle A(\tau) \rangle &= \text{Tr} \rho \hat{A}_H(\tau) \end{aligned} \quad (5.5)$$

We note that for two local observables A_1, A_2 the linear combinations $A = \alpha_1 A_1 + \alpha_2 A_2$ are also local observables. The associated operators obey the same linear relations $\hat{A} = \alpha_1 \hat{A}_1 + \alpha_2 \hat{A}_2$, where $\hat{A}$ stands for $\hat{A}_\delta, \hat{A}_S$ or $\hat{A}_H$. The relation (5.5) is the appropriate formula to answer the above question. One may use the set of measurements of expectation values at $\tau = 0$ to gather information about ρ . Once ρ is determined with sufficient accuracy, the expectation values $\langle A(\tau) \rangle$ can be computed. Of course, this needs a computation of the explicit form of the Heisenberg operator $\hat{A}_H(\tau)$.

It is instructive to observe that some simple local observables have an essentially τ -independent operator representation in the Schrödinger picture. This is easily seen for observables $A(\tau)$ which depend only on the variable $\varphi(\tau)$. The mapping reads

$$A(\tau) = f(\varphi(\tau)) \rightarrow \hat{A}_S(\tau) = f(\hat{Q}) \quad (5.6)$$

For the derivative operator $(\partial_\tau \varphi(\tau))^2$ a simple representation obtains in the limit $\epsilon \rightarrow 0$

$$\begin{aligned}
A(\tau) &= (\partial_\tau \varphi(\tau))^2 \rightarrow \\
\hat{A}_S(\tau) &= \frac{1}{\epsilon} \left(\frac{1}{Z(\tau + \epsilon) + Z(\tau)} + \frac{1}{Z(\tau - \epsilon) + Z(\tau)} \right) \\
&\quad - \hat{P}^2 \left(\frac{2}{(Z(\tau + \epsilon) + Z(\tau))^2} + \frac{2}{(Z(\tau - \epsilon) + Z(\tau))^2} \right) \\
&\quad - 2 \left\{ \frac{2}{(Z(\tau + \epsilon) + Z(\tau))^2} e^{\frac{\epsilon}{2} V(\hat{Q}, \tau)} \left[\hat{P}^2, e^{-\frac{\epsilon}{2} V(\hat{Q}, \tau)} \right] \right. \\
&\quad \left. - \frac{2}{(Z(\tau - \epsilon) + Z(\tau))^2} \left[\hat{P}^2, e^{-\frac{\epsilon}{2} V(\hat{Q}, \tau)} \right] e^{\frac{\epsilon}{2} V(\hat{Q}, \tau)} \right\} \quad (5.7)
\end{aligned}$$

The last term is $\sim \epsilon^2$ and for constant Z and $\epsilon \rightarrow 0$ one has

$$(\partial_\tau \varphi(\tau))^2 \rightarrow 1/(\epsilon Z) - \hat{P}^2/Z^2 \quad (5.8)$$

We observe that the r.h.s. of eq. (5.8) differs from $-\hat{P}^2/Z^2$ by a constant which diverges for $\epsilon \rightarrow 0$. This reconciles the positivity of $(\partial_\tau \varphi)^2$ with positive³ expectation values of $\hat{P}^2$.

For the transition to the Heisenberg picture the explicit definitions (3.3) and (3.10) yield

$$\hat{Q}\hat{U}(\tau) = \varphi(\tau)\hat{U}(\tau), \quad \hat{U}(\tau)\hat{Q} = \varphi(0)\hat{U}(\tau) \quad (5.9)$$

and we note that the evolution operator does not commute with $\hat{Q}$,

$$[\hat{Q}, \hat{U}(\tau)] = (\varphi(\tau) - \varphi(0))\hat{U}(\tau) = (\varphi_2 - \varphi_1)\hat{U}(\varphi_2, \varphi_1; \tau, 0) \quad (5.10)$$

Observables depending only on one variable $\varphi(\tau)$ have the Heisenberg representation (cf. eq(5.6))

$$A(\tau) = f(\varphi(\tau)) \rightarrow \hat{A}_H(\tau) = \hat{U}^{-1}(\tau)f(\hat{Q})\hat{U}(\tau) = f(\hat{Q}(\tau)) \quad (5.11)$$

Here we have used the definition

$$\hat{Q}(\tau) = \hat{U}^{-1}(\tau)\hat{Q}\hat{U}(\tau) \quad (5.12)$$

More generally, one finds for products of functions depending on the variables $\varphi(\tau_1), \varphi(\tau_2) \dots \varphi(\tau_n)$ with $\tau_1 < \tau_2 < \dots \tau_n$ the Heisenberg operator

$$\begin{aligned}
A(\tau_1, \dots \tau_n) &= f_1(\varphi(\tau_1))f_2(\varphi(\tau_2)) \dots f_n(\varphi(\tau_n)) \longrightarrow \\
\hat{A}_H(\tau) &= \hat{U}^{-1}(\tau_n)f_n(\hat{Q})\hat{U}(\tau_n, \tau_{n-1}) \dots \hat{U}(\tau_3, \tau_2)f_2(\hat{Q})\hat{U}(\tau_2, \tau_1)f_1(\hat{Q})\hat{U}(\tau_1) \\
&= f_n(\hat{Q}(\tau_n)) \dots f_2(\hat{Q}(\tau_2))f_1(\hat{Q}(\tau_1)) \quad (5.13)
\end{aligned}$$

³Note that $\hat{P}^2$ is not *necessarily* positive as long as $|\psi(\tau)\rangle$ and $\{\psi(\tau)|$ are unrelated.

This important relation follows directly from the definitions (2.10), (5.1), (5.5). We observe that $\hat{A}_H$ depends on the variables τ_i which are the arguments of A but shows no dependence on the reference point τ . (Only $\hat{A}_\delta$ and $\hat{A}_S$ depend on τ .)

We can use eq. (5.13) to find easily the Heisenberg operators for observables involving “derivatives”, e.g.

$$\begin{aligned}
A &= \tilde{\partial}_\tau \varphi(\tau_1) = \frac{1}{2\epsilon} (\varphi(\tau_1 + \epsilon) - \varphi(\tau_1 - \epsilon)) \\
\hat{A}_H &= \frac{1}{2\epsilon} \{ \hat{U}^{-1}(\tau_1 + \epsilon) \hat{Q} \hat{U}(\tau_1 + \epsilon) - \hat{U}^{-1}(\tau_1 - \epsilon) \hat{Q} \hat{U}(\tau_1 - \epsilon) \} \\
&= \frac{1}{2\epsilon} \hat{U}^{-1}(\tau_1) \{ e^{\epsilon \hat{H}(\tau_1 + \frac{\epsilon}{2})} \hat{Q} e^{-\epsilon \hat{H}(\tau_1 + \frac{\epsilon}{2})} - e^{-\epsilon \hat{H}(\tau_1 - \frac{\epsilon}{2})} \hat{Q} e^{\epsilon \hat{H}(\tau_1 - \frac{\epsilon}{2})} \} \hat{U}(\tau_1) \\
&= \hat{U}^{-1}(\tau_1) [\hat{H}(\tau_1), \hat{Q}] \hat{U}(\tau_1) + O(\epsilon) \\
&= \frac{1}{2Z(\tau_1)} \hat{U}^{-1}(\tau_1) [\hat{P}^2, \hat{Q}] \hat{U}(\tau_1) + O(\epsilon) \\
&= -\frac{1}{Z(\tau_1)} \hat{U}^{-1}(\tau_1) \hat{R} \hat{U}(\tau_1) + O(\epsilon) = -\frac{1}{Z(\tau_1)} \hat{R}(\tau_1) + O(\epsilon)
\end{aligned} \tag{5.14}$$

where we have assumed that $\hat{H}$ is a smooth function of τ . Here $\hat{R}$ is defined by

$$\hat{R}(\varphi_2, \varphi_1) = \delta(\varphi_2 - \varphi_1) \frac{\partial}{\partial \varphi_1}, \quad \hat{R}^2 = -\hat{P}^2 \tag{5.15}$$

and we use, similar to eq. (5.12), the definitions

$$\hat{R}(\tau) = \hat{U}^{-1}(\tau) \hat{R} \hat{U}(\tau), \quad \hat{P}^2(\tau) = \hat{U}^{-1}(\tau) \hat{P}^2 \hat{U}(\tau) \tag{5.16}$$

Two different definitions of derivatives can lead to the same operator $\hat{A}_H$. An example is the observable

$$A = \partial_\tau^> \varphi(\tau_1) = \frac{1}{\epsilon} (\varphi(\tau_1 + \epsilon) - \varphi(\tau_1)) \tag{5.17}$$

Up to terms of order ϵ the associated Heisenberg operator is again given by $\hat{A}_H = -Z(\tau_1)^{-1} \hat{U}^{-1}(\tau_1) \hat{R} \hat{U}(\tau_1)$ and therefore the same as for $\tilde{\partial}_\tau \varphi(\tau_1)$ (eq. (5.14)). Applying the same procedure to the squared derivative observable (2.4) yields

$$\begin{aligned}
A &= (\partial_\tau \varphi)^2(\tau_1) \longrightarrow \\
\hat{A}_H &= \frac{1}{2\epsilon^2} \hat{U}^{-1}(\tau_1) \{ 2\epsilon [\hat{Q}, [\hat{H}, \hat{Q}]] + 2\epsilon^2 [\hat{Q}, \hat{H}] [\hat{Q}, \hat{H}] \} \hat{U}(\tau_1) \\
&= \frac{1}{\epsilon Z} - \frac{1}{Z^2} \hat{U}^{-1}(\tau_1) \hat{P}^2 \hat{U}(\tau_1) = \frac{1}{\epsilon Z} - \frac{1}{Z^2} \hat{P}^2(\tau_1)
\end{aligned} \tag{5.18}$$

where we have assumed for simplicity a τ -independent Hamiltonian $\hat{H}$. This is in agreement with the direct evaluation (5.8). It is remarkable that this operator differs

from the square of the Heisenberg operator associated to $\tilde{\partial}_\tau \varphi(\tau_1)$ by a constant which diverges for $\epsilon \rightarrow 0$. Indeed, one finds

$$\begin{aligned} A &= (\tilde{\partial}_\tau \varphi(\tau_1))^2 \rightarrow \hat{A}_H = \frac{1}{2\epsilon Z} - \frac{1}{Z^2} \hat{P}^2(\tau_1) \\ A &= (\partial_\tau^> \varphi(\tau_1))^2 \rightarrow \hat{A}_H = \frac{1}{\epsilon Z} - \frac{1}{Z^2} \hat{P}^2(\tau_1) \end{aligned} \quad (5.19)$$

Incidentally, these relations can be used in order to find an observable whose Schrödinger operator is given by $\hat{P}^2$ and Heisenberg operator by $\hat{P}^2(\tau)$:

$$\begin{aligned} A_{P^2}(\tau_1) &= Z^2 (\partial_\tau^> \varphi(\tau_1))^2 - 2Z^2 (\tilde{\partial}_\tau \varphi(\tau_1))^2 \\ &= \frac{Z^2}{\epsilon^2} \{ \varphi^2(\tau_1) + \varphi(\tau_1 + \epsilon) \varphi(\tau_1 - \epsilon) - \varphi(\tau_1 + \epsilon) \varphi(\tau_1) - \varphi(\tau_1) \varphi(\tau_1 - \epsilon) \} \\ &\rightarrow \hat{A}_H = \hat{P}^2(\tau_1) \end{aligned} \quad (5.20)$$

As a last instructive example for the association between observables and Heisenberg operators we consider (for $\epsilon \rightarrow 0$)

$$\begin{aligned} A &= \partial_\tau^> \varphi(\tau_1) \varphi(\tau_2) \longrightarrow \\ \hat{A}_H &= \begin{cases} -Z^{-1} \hat{R}(\tau_1) \hat{Q}(\tau_2) & \text{for } \tau_1 \geq \tau_2 + \epsilon \\ -Z^{-1} \hat{Q}(\tau_2) \hat{R}(\tau_1) & \text{for } \tau_1 \leq \tau_2 - \epsilon \end{cases} \end{aligned} \quad (5.21)$$

The place where $\hat{R}$ and $\hat{Q}$ appear depends on the ordering of τ_1 and τ_2 . One obtains the same Heisenberg operator (5.21) if one replaces $\partial_\tau^> \varphi(\tau_1)$ by $\tilde{\partial}_\tau \varphi(\tau_1)$. On the other hand, for equal locations $\tau_2 = \tau_1$ the two observables correspond to different Heisenberg operators

$$\begin{aligned} \varphi(\tau_1) \partial_\tau^> \varphi(\tau_1) &\rightarrow -Z^{-1} \hat{U}^{-1}(\tau_1) \hat{R} \hat{Q} \hat{U}(\tau_1) \\ \varphi(\tau_1) \tilde{\partial}_\tau \varphi(\tau_1) &\rightarrow -\frac{1}{2} Z^{-1} \hat{U}^{-1}(\tau_1) (\hat{R} \hat{Q} + \hat{Q} \hat{R}) \hat{U}(\tau_1) \\ &= -Z^{-1} \hat{U}^{-1}(\tau_1) \hat{R} \hat{Q} \hat{U}(\tau_1) + \frac{1}{2} Z^{-1} \end{aligned} \quad (5.22)$$

One concludes that for arbitrary probability distributions $p_<$ and $p_>$ (or states $\{|\psi\rangle$ and $|\psi\rangle\}$) the expectation values $\langle \partial_\tau^> \varphi(\tau_1) \varphi(\tau_1) \rangle$ and $\langle \tilde{\partial}_\tau \varphi(\tau_1) \varphi(\tau_1) \rangle$ differ.

Eqs. (5.19) and (5.22) teach us that the product of derivative observables with other observables can be ambiguous in the sense that the associated operator and expectation value depends very sensitively on the precise definition of the derivative. This ambiguity of the derivative observables in the continuum limit is an unpleasant feature for the formulation of correlation functions. In the next sections we will see how this problem is connected with the concept of quantum correlations. We will argue that the ambiguity in the classical correlation may be the basic ingredient why a description of our world in terms of quantum statistics is superior to the use of classical correlation functions.

6 Correlation functions in incomplete classical statistics

A basic concept for any statistical description are correlation functions for a number of observables $A_1[\varphi], A_2[\varphi], \dots$. In particular, a two-point function is given by the expectation value of an associative product of two observables $A_1[\varphi]$ and $A_2[\varphi]$. For local observables A_1, A_2 the product should again be a local observable which must be defined uniquely in terms of the definitions of A_1 and A_2 . This requirement, however, does not fix the definition of the correlation uniquely. The standard “classical product”, i.e. the simple multiplication of the functionals $A_1[\varphi] \cdot A_2[\varphi]$ (in the same sense as the “pointwise” multiplication of functions) fulfills the general requirements⁴ for a correlation function. Other definitions can be conceived as well. In this section we will introduce a quantum correlation which equals the classical (“pointwise”) correlation only for τ -ordered non-overlapping observables. In contrast, for two local observables with overlapping support we will find important differences between the quantum and classical correlation. In particular, we will discover the effects of the non-commutativity characteristic for quantum mechanics.

Incomplete statistics draws our attention to an important issue in the formulation of meaningful correlation functions. Consider the two versions of the derivative observable $\tilde{\partial}_\tau \varphi$ and $\partial_\tau^> \varphi$ defined by eqs. (5.14) and (5.15), respectively. In the continuum limit ($\epsilon \rightarrow 0$) they are represented by the same operator $\hat{A}_H$. In consequence, both definitions lead to the same expectation value for any state $|\psi\rangle, \{\psi\}$. The two versions of derivative observables cannot be distinguished by any measurement and should therefore be identified. On the other hand, the classical products $\tilde{\partial}_\tau \varphi(\tau_1) \cdot \tilde{\partial}_\tau \varphi(\tau_2)$ and $\partial_\tau^> \varphi(\tau_1) \cdot \partial_\tau^> \varphi(\tau_2)$ are represented by different operators for $\tau_1 = \tau_2$, as can be seen from eq. (5.19). (See also eq. (5.22) for a similar argument involving $\partial_\tau \varphi(\tau_1) \cdot \varphi(\tau_2)$.) This means that the two versions of derivative observables lead to different classical correlation functions! Obviously, this situation is unsatisfactory since for $\epsilon \rightarrow 0$ no difference between the two versions could be “measured” for the observables themselves. We find this disease unacceptable for a meaningful correlation and require as a criterion for a meaningful correlation function that two observables which have the same expectation values for all (arbitrary) probability distributions should also have identical correlation functions. We have shown that two observables which are represented by the same Heisenberg operator have indeed the same expectation values for all possible probability distributions and should therefore be considered as equivalent. They should therefore lead to indistinguishable correlation functions.

As we have already established, the two derivative observables $A_1 = \tilde{\partial}_\tau \varphi(\tau)$ and $A_2 = \partial_\tau^> \varphi(\tau)$ are indistinguishable in the continuum limit, whereas their classical correlations are not. We may therefore conclude⁵ that the classical correlation

⁴This holds provided that the product results in a meaningful observable with finite expectation value.

⁵We will discuss in the next section to what extent this problem may be cured by a restriction

$A_1 \cdot A_2$ is not a meaningful correlation function. In this section we propose the use of a different correlation based on a quantum product $A_1 \circ A_2$. By construction, this correlation will always obey our criterion of “robustness” with respect to the precise choice of the observables. It should therefore be considered as an interesting alternative to the classical correlation. At this place we only note that the “robustness problem” is not necessarily connected to the continuum limit. The mismatch between indistinguishable observables and distinguishable “classical” correlations can appear quite generally also for $\epsilon > 0$.

Our formulation of a quantum correlation will be based on the concepts of equivalent observables and products defined for equivalence classes. In fact, the mapping $A(\tau) \rightarrow \hat{A}_H(\tau)$ is not necessarily invertible on the space of all observables $A(\tau)$. This follows from the simple observation that already the map (2.10) contains integrations. Two different integrands (observables) could lead to the same value of the integral (operator) for arbitrary fixed boundary values $\varphi(\tau - \frac{\delta}{2})$, $\varphi(\tau + \frac{\delta}{2})$. It is therefore possible that two different observables $A_a(\tau)$ and $A_b(\tau)$ can be mapped into the same Heisenberg operator $\hat{A}_H(\tau)$. Since the expectation values can be computed from $\hat{A}_H(\tau)$ and ρ only, no distinction between $\langle A_a \rangle$ and $\langle A_b \rangle$ can then be made for arbitrary ρ . All local observables $A(\tau)$ which correspond to the same operator $\hat{A}_H(\tau)$ are equivalent.

We are interested in structures that only depend on the equivalence classes of observables. Addition of two observables and multiplication with a scalar can simply be carried over to the operators. This is not necessarily the case, however, for the (pointwise) multiplication of two observables. If $A_a(\tau)$ and $A_b(\tau)$ are both mapped into $\hat{A}_H(\tau)$ and a third observable $B(\tau)$ corresponds to $\hat{B}_H(\tau)$, the products $A_a \cdot B$ and $A_b \cdot B$ may nevertheless be represented by different operators (cf. eq. (5.20), (5.22)). It is then easy to construct states where $\langle A_a B \rangle \neq \langle A_b B \rangle$ and the pointwise product does not depend only on the equivalence class.

On the other hand, the (matrix) product of two operators $\hat{A}_H \hat{B}_H$ obviously refers only to the equivalence class. It can be implemented on the level of observables by defining a unique “standard representative” of the equivalence class as

$$\bar{A}[\varphi, \tau] = F[\hat{A}_H(\tau)] \quad (6.1)$$

Using the mapping $A[\tau] \rightarrow \hat{A}_H(\tau)$ (2.10), (5.1), (5.5), we define the quantum product of two observables as

$$A(\varphi, \tau) \circ B(\varphi, \tau) = F[\hat{A}_H(\tau) \hat{B}_H(\tau)] \equiv (A \circ B)[\varphi, \tau] \quad (6.2)$$

This product is associative, but not commutative. (By definition, the operator associated to the observable $(A \circ B)(\varphi, \tau)$ is $\hat{A}_H(\tau) \hat{B}_H(\tau)$ and the product $A \circ B$ is isomorphic to the “matrix multiplication” $\hat{A} \hat{B}$ if restricted to the subspace of operators $\bar{A} = F[\hat{A}]$, $\bar{B} = F[\hat{B}]$.) The correlations (e.g. expectation values of products of observables) formed with the product $\circ$ reflect the non-commutative

to better “smoothened” observables.

structure of quantum mechanics. This justifies the name “quantum correlations”. Nevertheless, we emphasize that the “quantum product” $\circ$ can also be viewed as just a particular structure among “classical observables”.

The definition of the quantum product is unique on the level of operators. On the level of the classical observables, it is, however, not yet fixed uniquely by eq. (6.2). The precise definition obviously depends on the choice of a standard representation $F[\hat{A}_H(\tau)]$ for the equivalence class of observables represented by $\hat{A}_H$. We will choose a linear map $F[\alpha_1\hat{A}_{H,1} + \alpha_2\hat{A}_{H,2}] = \alpha_1F[\hat{A}_{H,1}] + \alpha_2F[\hat{A}_{H,2}]$ with the property that it inverts the relation (5.13). For “time-ordered” $\tau_1 < \tau_2 < \dots \tau_n$ the map F should then obey

$$F[f_n(\hat{Q}(\tau_n)) \dots f_2(\hat{Q}(\tau_2))f_1(\hat{Q}(\tau_1))] = f_1(\varphi(\tau_1))f_2(\varphi(\tau_2)) \dots f_n(\varphi(\tau_n)). \quad (6.3)$$

It is easy to see how this choice exhibits directly the noncommutative property of the quantum product between two observables. As an example let us consider the two observables $\varphi(\tau_1)$ and $\varphi(\tau_2)$ with $\tau_1 < \tau_2$. The quantum product or quantum correlation depends on the ordering

$$\begin{aligned} \varphi(\tau_2) \circ \varphi(\tau_1) &= \varphi(\tau_2)\varphi(\tau_1) \\ \varphi(\tau_1) \circ \varphi(\tau_2) &= \varphi(\tau_2)\varphi(\tau_1) + F[[\hat{Q}(\tau_1), \hat{Q}(\tau_2)]] \end{aligned} \quad (6.4)$$

The noncommutative property of the quantum product for these operators is directly related to the commutator

$$\begin{aligned} [\hat{Q}(\tau_1), \hat{Q}(\tau_2)] &= \hat{U}^{-1}(\tau_1)\hat{Q}\hat{U}(\tau_1, \tau_2)\hat{Q}\hat{U}(\tau_2) \\ &\quad - \hat{U}^{-1}(\tau_2)\hat{Q}\hat{U}(\tau_2, \tau_1)\hat{Q}\hat{U}(\tau_1) \end{aligned} \quad (6.5)$$

Only for time-ordered arguments the quantum correlation coincides with the classical correlation. The nonvanishing commutator $\varphi(\tau_1) \circ \varphi(\tau_2) - \varphi(\tau_2) \circ \varphi(\tau_1)$ could presumably play a crucial role in the discussion of Bell’s inequalities [4] once our system is extended to a setting with more than one particle.⁶

The map F can easily be extended to operators involving derivatives of φ . We concentrate here for simplicity on a translation invariant probability distribution in the local region with constant $Z(\tau) = Z$. The mappings (with $\tau_2 \geq \tau_1 + \epsilon$)

$$\begin{aligned} F(\hat{R}(\tau)) &= -Z\partial_\tau^>\varphi(\tau) \\ F(\hat{R}(\tau)\hat{Q}(\tau)) &= -Z\varphi(\tau)\partial_\tau^>\varphi(\tau) \\ F(\hat{R}(\tau_2)\hat{R}(\tau_1)) &= Z^2\partial_\tau^>\varphi(\tau_2)\partial_\tau^>\varphi(\tau_1) \end{aligned} \quad (6.6)$$

are compatible with eq. (6.3). This can be seen by noting that the τ -evolution of $\hat{Q}(\tau)$ according to eq. (5.12) implies for $\epsilon \rightarrow 0$ the simple relation

$$\partial_\tau\hat{Q}(\tau) = [\hat{H}, \hat{Q}(\tau)] = -Z^{-1}\hat{R}(\tau) \quad (6.7)$$

⁶Note that Bell’s inequalities are based on the definition of the correlation function by “pointwise multiplication”.

A similar construction (note $[\hat{Q}(\tau + \epsilon), \hat{Q}(\tau)] = -\epsilon/Z$) leads to

$$F(\hat{R}^2(\tau)) = Z^2(\partial_\tau^>\varphi(\tau))^2 - Z/\epsilon \quad (6.8)$$

and we infer that the quantum product of derivative observables at equal sites differs from the pointwise product

$$\partial_\tau^>\varphi(\tau) \circ \partial_\tau^>\varphi(\tau) = (\partial_\tau^>\varphi(\tau))^2 - 1/(\epsilon Z) \quad (6.9)$$

From the relations (6.4) and (6.9) it has become clear that the difference between the quantum product and the “pointwise” classical product of two observables is related to their τ -ordering and “overlap”. Let us define that two observables $A_1[\varphi]$ and $A_2[\varphi]$ overlap if they depend on variables $\varphi(\tau)$ lying in two overlapping τ -ranges $\mathcal{R}_1$ and $\mathcal{R}_2$. (Here two ranges do not overlap if all τ in $\mathcal{R}_1$ obey $\tau \leq \tau_0$ whereas for $\mathcal{R}_2$ one has $\tau \geq \tau_0$, or vice versa. This implies that non-overlapping observables can depend on at most one common variable $\varphi(\tau_0)$.) With this definition the quantum product is equal to the classical product if the observables do not overlap and are τ -ordered (in the sense that larger τ are on the left side).

In conclusion, we have established a one-to-one correspondence between classical correlations $\varphi(\tau_2)\varphi(\tau_1)$ and the product of Heisenberg operators $\hat{Q}(\tau_2)\hat{Q}(\tau_1)$ provided that the τ -ordering $\tau_2 \geq \tau_1$ is respected. This extends to observables that can be written as sums or integrals over $\varphi(\tau)$ (as, for example, derivative observables) provided the τ -ordering and non-overlapping properties are respected. For well separated observables no distinction between a quantum and classical τ -ordered correlation function would be needed. In particular, this holds also for “smoothened” observables A_i that involve (weighted) averages over $\varphi(\tau)$ in a range $\mathcal{R}_i$ around τ_i . Decreasing the distance between τ_2 and τ_1 , the new features of the quantum product $A_1(\tau_2) \circ A_1(\tau_1)$ show up only once the distance becomes small enough so that the two ranges $\mathcal{R}_1$ and $\mathcal{R}_2$ start to overlap. In an extreme form the difference between quantum and classical correlations becomes apparent for derivative observables at the same location. Quite generally, the difference between the quantum and classical product is seen most easily on the level of the associated operators

$$\begin{aligned} A_1 \circ A_2 &\rightarrow \hat{A}_1 \hat{A}_2 \\ A_1 \cdot A_2 &\rightarrow T(\hat{A}_1 \hat{A}_2) \end{aligned} \quad (6.10)$$

Here T denotes the operation of τ -ordering. The τ -ordered operator product is commutative $T(\hat{A}_1 \hat{A}_2) = T(\hat{A}_2 \hat{A}_1)$ and associative $T(T(\hat{A}_1 \hat{A}_2) \hat{A}_3) = T(\hat{A}_1 T(\hat{A}_2 \hat{A}_3)) \equiv T(\hat{A}_1 \hat{A}_2 \hat{A}_3)$. As we have seen in the discussion of the derivative observables, it lacks, however, the general property of robustness with respect to the precise definition of the observables. This contrasts with the non-commutative product $\hat{A}_1 \hat{A}_2$.

7 Complete classical statistics, irrelevant and inaccessible information

Our discussion of incomplete classical statistics may perhaps have led to the impression that the quantum mechanical properties are somehow related to the missing information. This is by no means the case! In fact, our investigation of the consequences of incomplete information about the probability distribution was useful in order to focus the attention on the question which information is really necessary to compute the expectation values of local observables. We can now turn back to standard “complete” classical statistics where a given probability distribution $p[\varphi(\tau)]$ is assumed to be known. We concentrate here on a general class of probability distributions which can be factorized in the form $p = p_>p_0p_<$ according to eq. (1.1) – it may be called “factorizable” or “ F -statistics”. For example, all systems which have only local and next-neighbor interactions are of this form. Within F -statistics the states remain defined according to eq. (2.9).

We emphasize that any additional information contained in $p[\varphi]$ which goes beyond the local distribution $p_0[\varphi]$ and the states $|\psi\rangle$ and $\langle\psi|$ does not change a iota in the expectation values of local observables and their correlations! The additional information is simply *irrelevant* for the computation of local expectation values. A given probability distribution specifies $p_<$ and $p_>$ uniquely. This determines $|\psi\rangle$ and $\langle\psi|$ and we can then continue with the preceding discussion in order to calculate the expectation values of local observables. The precise form of $p_<$ and $p_>$ which has led to the given states plays no role in this computation.

Since all information contained in $p_<$ and $p_>$ beyond the states $|\psi\rangle$ and $\langle\psi|$ is irrelevant for local expectation values, it is also *inaccessible* by any local measurements. In fact, even the most precise measurements of expectation values and correlation functions for arbitrarily many local observables could at best lead to a reconstruction of the states $|\psi\rangle$ and $\langle\psi|$. This sheds new light on the notion of “incompleteness” of the statistical information discussed in this note. In fact, within F -statistics the “incomplete” information contained in the states $|\psi\rangle$ and $\langle\psi|$ constitutes the most complete information that can possibly be achieved by local measurements! Since any real measurement is local in time and space all assumptions about information beyond the states concern irrelevant and inaccessible information and cannot be verified by observation!

Particularly interesting in the spirit of the completeness question asked by Einstein is the following situation: Suppose first that an observer has no information about the “exterior region” beyond the one contained in the state vectors. Nevertheless, his information is sufficient in order to compute the expectation values and correlations of all local observables. What happens, however, if additional information about the probability distribution in the exterior region becomes available? Our observer could now perhaps compute additional expectation values for observables with support outside the local region. As far as the local observables are concerned, however, the additional information is completely irrelevant. In this sense quantum

mechanics is complete as far as local observables are concerned.

These simple remarks have striking consequences: all the quantum mechanical features found in our discussion of incomplete classical statistics are genuine properties of *all* classical statistical systems whose probability distribution can be factorized. This concerns, in particular, the expectation values of the observables $\varphi^p(\tau)$. For an arbitrary τ -translation invariant and factorizable probability distribution all $\langle \varphi^p(\tau) \rangle$ will be independent of τ unless the translation symmetry is spontaneously broken. We infer that only the discrete stationary values of $\langle \hat{Q}^p \rangle$ of the quantum-mechanical system with the appropriate $\hat{H}$ are possible solutions. We see how “quantum mechanical” information can be used directly for a discussion of classical statistical properties in a wide class of systems.

Another possibility concerns the use of the uncertainty relation which is also present in classical F -statistics. Assuming for simplicity a discrete symmetry $\varphi \rightarrow -\varphi$ with $\langle \varphi \rangle = 0$ this yields the inequality (cf. the definition (5.20))

$$\langle \varphi^2(\tau) \rangle \langle A_{P^2}(\tau) \rangle = \langle \hat{Q}^2 \rangle \langle \hat{P}^2 \rangle \geq 1/4 \quad (7.1)$$

(Throughout this paper we use conventions for the units of φ and τ where $\hbar \equiv 1$.) It is not obvious how such an inequality for the derivative-like observable A_{P^2} (5.20) would have been found without using the simple commutator relations of quantum-mechanical operators. The simple fact that quantum-mechanical information can be used in practice to establish properties of expectation values in a standard classical statistical system demonstrates in a simple way that quantum-mechanical features are indeed genuine properties of classical statistical systems.

The quantum correlation between two observables can be formulated in standard classical statistics as well. Together with the “classical” correlation function we have thus more than one candidate for the definition of a correlation function in a given classical statistical system⁷. The final decision which correlation is relevant in practice depends on the question which type of correlation can typically be measured by experiment. We believe that a certain “robustness” of the correlation with respect to the precise choice of the observable plays an important role in this respect.

We have argued in the preceding section that the classical correlation may be disfavored by the lack of robustness for local correlations of local derivative operators. The observed disease of the classical correlation gets weakened, however, once we consider the average of a correlation $\langle A(\tau_1)B(\tau_2) \rangle$ over a certain range in $|\tau_2 - \tau_1|$. With the mappings from observables to Heisenberg operators (for $\epsilon \rightarrow 0$, $Z(\tau) = Z$)

$$\begin{aligned} \partial_\tau^> \varphi(\tau + \epsilon) \partial_\tau^> \varphi(\tau) &\rightarrow \frac{1}{Z^2} \hat{R}(\tau + \epsilon) \hat{R}(\tau) = -\frac{1}{Z^2} \hat{P}^2(\tau) \\ \tilde{\partial}_\tau \varphi(\tau + \epsilon) \tilde{\partial}_\tau \varphi(\tau) &= \frac{1}{4\epsilon Z} - \frac{1}{Z^2} \hat{P}^2 \end{aligned} \quad (7.2)$$

⁷Actually, the same situation would arise had we started from a quantum-mechanical system: one has to decide if the noncommuting operator product $\hat{A}\hat{B}$ or rather the commuting τ -ordered product $T(\hat{A}\hat{B})$ should be used for the definition of the correlation.

we see that the difference between the two versions of derivative operators vanishes for an average of classical correlations which is defined by

$$\frac{1}{3} \sum_{m=-1}^1 < \partial_\tau \varphi(\tau + m\epsilon) \partial_\tau \varphi(\tau) > \quad (7.3)$$

Similarly, for $\Delta = (2n+1)\epsilon$ we may define averaged observables

$$A^\Delta(\tau) = \frac{1}{2n+1} \sum_{m=-n}^n A(\tau + \epsilon n) \quad (7.4)$$

and we find for the difference of the square of the two versions of derivative operators

$$(\partial_\tau^>\varphi)^\Delta(\tau)(\partial_\tau^>\varphi)^\Delta(\tau) - (\tilde{\partial}_\tau^\Delta\varphi)^\Delta(\tau)(\tilde{\partial}_\tau^\Delta\varphi)^\Delta(\tau) \rightarrow \frac{\epsilon}{2\Delta^2} \quad (7.5)$$

For fixed Δ this vanishes for $\epsilon \rightarrow 0$ and one may incorrectly conclude that the problem of the lack of robustness for classical correlation functions disappears for smoothened or averaged observables.

That this is not the case can be realized by the comparison of two other smoothened derivative observables

$$\begin{aligned} \partial_\tau^{\Delta>}\varphi(\tau) &= \frac{\varphi(\tau + \Delta) - \varphi(\tau)}{\Delta}, \\ \tilde{\partial}_\tau^\Delta\varphi(\tau) &= \frac{\varphi(\tau + \Delta) - \varphi(\tau - \Delta)}{2\Delta} \end{aligned} \quad (7.6)$$

It is straightforward to verify that both $(\partial_\tau^>\varphi)^\Delta$ and $(\tilde{\partial}_\tau^\Delta\varphi)^\Delta$ actually correspond to the observable $\tilde{\partial}_\tau^\Delta\varphi$ up to corrections $O(\epsilon)$ whereas for a discussion of the robustness problem on the scale Δ we should compare two versions of derivative observables that may differ in order $O(\Delta)$ as the two versions in eq. (7.6):

$$\begin{aligned} \partial_\tau^{\Delta>}\varphi(\tau) &\rightarrow -\frac{1}{Z}\hat{R}(\tau) + O(\Delta) \\ \tilde{\partial}_\tau^\Delta\varphi(\tau) &\rightarrow -\frac{1}{Z}\hat{R}(\tau) + O(\Delta) \end{aligned} \quad (7.7)$$

Nevertheless, the classical correlation function differs for the two choices by a term that does not vanish for $\Delta \rightarrow 0$: One finds, for $|\Delta < \hat{H}| \ll 1$ and $\tau_2 \geq \tau_1$:

$$\begin{aligned} &\partial_\tau^{\Delta>}\varphi(\tau_2)\partial_\tau^{\Delta>}\varphi(\tau_1) - \tilde{\partial}_\tau^\Delta\varphi(\tau_2)\tilde{\partial}_\tau^\Delta\varphi(\tau_1) \\ &\rightarrow \frac{\Delta - (\tau_2 - \tau_1)}{\Delta^2 Z} \theta(\Delta - (\tau_2 - \tau_1)) - \frac{\Delta - \frac{1}{2}(\tau_2 - \tau_1)}{2\Delta^2 Z} \theta(2\Delta - (\tau_2 - \tau_1)) \end{aligned} \quad (7.8)$$

In contrast to eq. (7.5) this difference survives in the continuum limit $\epsilon \rightarrow 0$. Furthermore, for $\tau_2 = \tau_1$ it “diverges” again for small Δ as $1/(2\Delta Z)$. We conclude

that the lack of robustness of the classical correlation remains present for averaged observables as well!

To be more precise, we will formulate the following “robustness criterion” for meaningful correlation functions: if two smoothened observables with support in a τ -range of the order $O(\Delta)$ have equal expectation values for all probability distributions up to corrections $\sim O(\Delta)$, then the respective correlation functions should also become identical up to corrections $\sim O(\Delta)$. (Here it is understood that $|\Delta < \hat{H} >| \ll 1$.) From eq. (7.8) we learn that the classical correlation does not meet this robustness criterion. For example, for τ_2 near τ_1 the difference between the correlation for the two versions of the derivative operator “diverges” $\sim \Delta^{-1}$. In fact, this disease extends to a whole range of $(\tau_2 - \tau_1) \sim O(\Delta)$. On the other hand, the quantum correlation obeys the robustness criterion by construction.

This discussion opens an interesting perspective: The difference between classical and quantum statistics seems to be a question of the appropriate definition of the correlation function. Simple arguments of robustness favor the choice of the quantum correlation! In a sense, the successful description of nature by quantum-mechanical operators and their products gives an “experimental indication” that quantum correlations should be used!

8 Superposition of states and density matrices

A fundamental principle of quantum mechanics is the superposition of states. It states that for two quantum states $|\psi^{(1)}\rangle$ and $|\psi^{(2)}\rangle$ the linear superposition $|\psi\rangle = \alpha_1|\psi^{(1)}\rangle + \alpha_2|\psi^{(2)}\rangle$ is again a possible quantum state. (Proper normalization is assumed.) Furthermore, for two density matrices $\rho^{(1)}, \rho^{(2)}$ the linear combination $\rho = w_1\rho^{(1)} + w_2\rho^{(2)}$, $w_1 + w_2 = 1$, is again a possible density matrix if it obeys the appropriate positivity conditions. We will show that these properties arise naturally in our context of incomplete classical statistics (or, more generally, F-statistics), at least for a restricted choice of α_1, α_2 .

Let us consider two probability distributions $p^{(1)}[\varphi] = \exp(-S^{(1)}[\varphi])$, $p^{(2)}[\varphi] = \exp(-S^{(2)}[\varphi])$ which differ only outside the local range $-\bar{\tau} < \tau < \bar{\tau}$, i.e.

$$\begin{aligned} S^{(i)}[\varphi] &= S_0^{(i)}[\varphi] + S_{>}^{(i)}[\varphi] + S_{<}^{(i)}[\varphi], \\ S_0^{(1)}[\varphi] &= S_0^{(2)}[\varphi] = S_0[\varphi] = \int_{-\bar{\tau}}^{\bar{\tau}} d\tau' \mathcal{L}(\tau'), \\ S_{>}^{(1)}[\varphi] &\neq S_{>}^{(2)}[\varphi], \quad S_{<}^{(1)}[\varphi] \neq S_{<}^{(2)}[\varphi] \end{aligned} \tag{8.1}$$

Within the local range the two probability distributions correspond to the same dynamics

$$\hat{H}^{(1)}(\tau) = \hat{H}^{(2)}(\tau), \quad \hat{U}^{(1)}(\tau) = \hat{U}^{(2)}(\tau) \tag{8.2}$$

whereas the states $|\psi^{(1)}\rangle$ and $|\psi^{(2)}\rangle$ differ. We will assume that both $S^{(1)}$ and $S^{(2)}$ are invariant under the reflection symmetry. As far as local observables are concerned the corresponding operators are the same for both situations. The only difference in the expectation values of observables between the two ensembles can be traced back to different state vectors at some reference point τ_0 or time t_0 . This setting reflects precisely the situation for two different quantum mechanical states $|\psi_1(t_0)\rangle \neq |\psi_2(t_0)\rangle$.

A superposition state corresponds to a new probability distribution given by $S[\varphi] = S_0[\varphi] + S_>[\varphi] + S_<[\varphi]$ with

$$\exp(-S_>[\varphi]) = \alpha_1 \exp(-S_>^{(1)}[\varphi]) + \alpha_2 \exp(-S_>^{(2)}[\varphi]) \quad (8.3)$$

Here the real coefficients α_1 and α_2 have to obey the condition that $\exp(-S[\varphi])$ is positive semidefinite for all φ and properly normalized. The definition (2.9) of the states as functional integrals implies directly

$$|\psi(\bar{\tau})\rangle = \alpha_1 |\psi^{(1)}(\bar{\tau})\rangle + \alpha_2 |\psi^{(2)}(\bar{\tau})\rangle \quad (8.4)$$

We observe that the superposition is compatible with the evolution such that for all τ in the interval $-\bar{\tau} < \tau < \bar{\tau}$ one finds $|\psi(\tau)\rangle = \alpha_1 |\psi^{(1)}(\tau)\rangle + \alpha_2 |\psi^{(2)}(\tau)\rangle$. We conclude that quantum mechanical superposition arises directly from the construction of the probability distribution (8.3). In particular, we emphasize the appearance of interference in the computation of expectation values of operators

$$\begin{aligned} \langle A \rangle = & \langle \psi | A^{(M)} | \psi \rangle = \alpha_1^2 \langle \psi^{(1)} | A^{(M)} | \psi^{(1)} \rangle + \alpha_2^2 \langle \psi^{(2)} | A^{(M)} | \psi^{(2)} \rangle \\ & + \alpha_1 \alpha_2 (\langle \psi^{(1)} | A^{(M)} | \psi^{(2)} \rangle + \langle \psi^{(2)} | A^{(M)} | \psi^{(1)} \rangle) \end{aligned} \quad (8.5)$$

The interference terms $\sim \alpha_1 \alpha_2$ are characteristic for “quantum statistics”. In our approach they are connected to the fact that the probability distribution corresponding to (8.3) cannot be written as a sum of two probability distributions! It should be emphasized that α_1 and α_2 may have opposite sign such that interference may indeed be destructive. Nevertheless, the condition stipulated after eq. (8.3) does not admit arbitrary superpositions. We note that the linear structure of the Schrödinger equation and the quantum mechanical time evolution arise directly from the basic construction. Nonlinear generalizations of quantum mechanics would have to modify this structure.

In distinction to the construction (8.3) we can also consider linear combinations of probability distributions

$$p[\varphi] = \exp(-S[\varphi]) = w_1 \exp(-S^{(1)}[\varphi]) + w_2 \exp(-S^{(2)}[\varphi]) \quad (8.6)$$

This alternative construction does not lead to interference and results in a linear combination of density matrices (cf. eq. (5.2))

$$\rho = w_1 \rho^{(1)} + w_2 \rho^{(2)} \quad (8.7)$$

with

$$\langle A \rangle = \text{Tr}(\hat{A}\rho) = w_1 \text{Tr}(\hat{A}\rho^{(1)}) + w_2 \text{Tr}(\hat{A}\rho^{(2)}) \quad (8.8)$$

The density matrix ρ (8.7) is not anymore the density matrix corresponding to a pure quantum mechanical state. In particular, the “pure state density matrices” $\rho^{(1)}, \rho^{(2)}$ obey

$$\text{Tr}(\rho^{(i)})^2 = \text{Tr}(\rho^{(i)}) = 1 \quad (8.9)$$

whereas (for $w_i \neq 0$)

$$\begin{aligned} \text{Tr}\rho^2 &= w_1^2 \text{Tr}(\rho^{(1)})^2 + w_2^2 \text{Tr}(\rho^{(2)})^2 + 2w_1w_2 \text{Tr}(\rho^{(1)}\rho^{(2)}) \\ &= 1 + 2w_1w_2(\text{Tr}(\rho^{(1)}\rho^{(2)}) - \text{Tr}\rho^{(1)}\text{Tr}\rho^{(2)}) \neq 1. \end{aligned} \quad (8.10)$$

We emphasize that the probability distribution (8.6) cannot be written in the product form (1.1) any more. The linear combination of density matrices therefore generalizes our original concept of incomplete statistics. This contrasts to the superposition of states (8.3) which preserves the product structure (1.1). For an extended concept of incomplete statistics with probability distributions of the type (8.6) new possibilities for τ -independent $\langle \varphi^p(\tau) \rangle$ arise, corresponding to stationary density matrices in quantum mechanics.

9 Conclusions and discussion

Within a simple example of classical statistics for coupled unharmonic oscillators on a chain we have formulated a description in terms of states and operators in analogy to quantum mechanics. The state vectors and the operators can be expressed in terms of classical functional integrals. Typical quantum mechanical results like the relations between the expectation values in stationary states or the uncertainty relation can be taken over to the classical system. Our procedure inverts the construction of the Euclidean path integral for a quantum mechanical system in the ground state or thermal state, with a generalization to a wider class of states.

The introduction of “quantum mechanical” operators associated to every local classical observable allows us to define equivalence classes of observables which cannot be distinguished by any measurement of their expectation values. We argue that the definition of the correlation function should be consistent with this equivalence structure. We require that indistinguishable observables must lead to the same correlation function. This leads to the introduction of a quantum correlation within the classical statistical setting. We point out that the quantum correlation constitutes a more robust definition of the correlation function with respect to the precise details of the definition of observables, both for classical and quantum statistical systems. The basic conceptual distinction between quantum statistics and classical statistics disappears in this respect. We have also seen the emergence of typical characteristics of quantum statistics like the superposition of states and interference. All this

points into the direction that it may be possible to understand the mysteries of the basics of quantum mechanics within a formulation of a classical statistical problem with infinitely many degrees of freedom.

What is missing is an understanding how the complex structure of quantum mechanics - the important phases - could be rooted in a possible classical statistical system. For our present system we discuss in the two appendices that a complex structure can be defined, if the system is invariant under a reflection symmetry. There we introduce a description in terms of normalized complex state vectors $|\psi(t)\rangle$ which depend on a real-time variable t . The time evolution is then given by the Schrödinger equation $i\partial_t|\psi\rangle = \hat{H}|\psi\rangle$ with a hermitean Hamilton operator (3.11). Expectation values of observables can be computed from associated operators in the usual way, e.g.

$$\langle \hat{Q}^2 \rangle = \langle \psi | \hat{Q}^2 | \psi \rangle \quad (9.1)$$

These operators obey the usual commutation relations, e.g.

$$[\hat{Q}, \hat{P}^2] = 2\hat{R} \quad , \quad [\hat{Q}, \hat{R}] = -1 \quad (9.2)$$

where $\hat{R}$ plays the role of $i\hat{P}$. As familiar in quantum mechanics we can equivalently use a Heisenberg or Schrödinger picture for the computation of the time evolution of expectation values of observables.

We observe, nevertheless, that at the present stage several insufficiencies remain on our way of understanding quantum mechanics from a classical statistical formulation. A major problem concerns the restriction that our description of states as functional integrals is restricted to those states $|\psi(t)\rangle$ whose analytical continuation $|\psi(\tau)\rangle$ for $t = -i\tau$ is a real positive function. Similarly, we have so far dealt only with linear superpositions $\alpha_1|\psi^{(1)}\rangle + \alpha_2|\psi^{(2)}\rangle$ with restricted real coefficients α_i such that the sum is again a real positive function. This situation seems deeply related to the issue of the complex structure and the important unsolved questions concerning the analytic continuation and the role of time.

The quantum mechanical features discussed in this note are present in all classical statistical systems. However, their conceptual meaning has become particularly clear in our formulation of incomplete classical statistics. The approach to incomplete statistics can be extended in various directions. First of all, incomplete statistical information does not necessarily occur in the form of missing information outside a local range. For example, the incompleteness of the information about the probability distribution can also concern the resolution within a given local interval. This problem probably takes a direction which is qualitatively very different from our development of quantum mechanics. Within the context used in this paper we may impose the additional restriction that only information which survives in the continuum limit $\epsilon \rightarrow 0$ is available. Otherwise stated, the resolution is not arbitrarily accurate. As discussed in the beginning of sect. 6 the different derivative observables become indistinguishable in such a version of incomplete statistics.

One can also investigate the consequences of abandoning certain of our assumptions. Without the reflection symmetry θ we obtain a description which remains

similar to quantum mechanics in many aspects. The Hamilton operator $\hat{H}$ does not remain hermitean, however, and the evolution operator $\hat{U}$ is not unitary any more (after analytic continuation). Without translation symmetry in τ we expect to find non-zero expectation values of antihermitean operators. We have not explored what happens if the interactions go beyond next-neighbor interactions. Finally, our approach can be extended to a collection of variables $\varphi_a(\tau)$. If we interpret a as a momentum label, we find a straightforward generalization to quantum field theory.

At this stage it is still open if the constructions presented here will find some practical use. Also the crucial question if quantum mechanics can indeed be formulated within a classical statistical system with infinitely many degrees of freedom has not yet found a definite answer. We hope that the finding of several structures characteristic for quantum mechanics within a classical statistical formulation will encourage the continued exploration in this direction.

Acknowledgment:

The author would like to thank M. Lüscher for useful discussions.

Appendix

A Reflection symmetry and complex structure

We have found several quantum-mechanical features in a classical statistical system, including the characteristic non-commutativity of operator products. In a sense, the quantum-mechanical and classical descriptions can be viewed as dual descriptions of one and the same system, distinguished by the choice of the correlation function. This raises the question if the other characteristic features of quantum mechanics, as the complex structure and the unitary time evolution, can also be found within classical statistics. In the two appendices we show that these structures may indeed be present on a formal level. So far we have not found, however, a fully consistent description of the quantum mechanical phases within our classical statistical setting.

Quantum mechanics has an intrinsic complex structure. This is manifest in the Schrödinger equation. In this section we establish how this complex structure emerges from a reflection symmetry in (incomplete) classical statistics. Let us consider the reflection⁸ at a given reference point $\tau = 0$, i.e.

$$\theta(\varphi(\tau)) = \varphi(-\tau), \quad \theta^2 = 1 \tag{A.1}$$

⁸Existence of the reflection for every τ requires that for every variable $\varphi(\tau)$ there is also a variable $\varphi(-\tau)$.

We concentrate on reflection-invariant probability distributions [9]

$$\begin{aligned}\theta(S[\varphi]) &= S[\varphi] \\ \theta(\mathcal{L}(\tau)) &= \mathcal{L}(-\tau) \\ \theta(S_{>}(\tau)) &= S_{<}(-\tau)\end{aligned}\tag{A.2}$$

for which $V_{\tau}(\varphi) = V_{-\tau}(\varphi)$, $Z_{\tau} = Z_{-\tau}$. The corresponding transformation properties of the states are

$$\theta|\psi(\tau)\rangle = |\psi(-\tau)\rangle, \quad \theta\{\psi(\tau)\} = \{\psi(-\tau)\}\tag{A.3}$$

We define the action of θ on matrices such that it also involves a transposition in the sense that $\theta(\{\psi_1|\hat{A}|\psi_2\}) = \theta|\psi_2\rangle\langle\psi_1|\theta\hat{A}$. Using the definitions of ρ , $\hat{U}$, $\hat{H}$ and the operators $\hat{A}$ one finds the relations

$$\begin{aligned}\theta\rho(\tau) &= \rho^T(-\tau), \quad \theta\hat{A}_{H,S,\delta}(\tau) = \hat{A}_{H,S,\delta}^{(R)T}(-\tau), \\ \theta\hat{H}(\tau) &= \hat{H}^T(-\tau), \quad \theta\hat{U}(\tau_2, \tau_1) = \hat{U}^T(-\tau_1, -\tau_2)\end{aligned}\tag{A.4}$$

Here $\hat{A}_{\delta}^{(R)}$ is related to $\hat{A}_{\delta}$ by replacing in eq. (2.10) for $A[\varphi(\tau + \eta)_{-\frac{\delta}{2} < \eta < \frac{\delta}{2}}]$ any $\varphi(\tau + \eta)$ by $\varphi(\tau - \eta)$, corresponding to a reflection of $A[\varphi, \tau]$ at τ . In consequence of the reflection symmetry of the probability distribution, the expectation value of any local observable must be equal to the one of the reflected observable

$$\theta \langle A(\tau) \rangle = \langle A^{(R)}(-\tau) \rangle = \langle A(\tau) \rangle\tag{A.5}$$

On the other hand, the reflection (A.1) acts on $|\psi\rangle$ only by a variable charge, i.e.

$$\theta|\psi(\varphi(\tau); \tau)\rangle = |\psi(\varphi(-\tau)); \tau\rangle = \{\psi(\varphi(-\tau)); -\tau|\tag{A.6}$$

Interpreting $|\psi\rangle$ and $\{\psi|\$ as functions of two variables φ and τ (without distinction to which τ the variable φ was originally associated) we can write

$$|\psi(-\tau)\rangle = \{\psi(\tau)|\tag{A.7}$$

and similarly $\theta(\rho(\tau)) = \rho(\tau)$ etc., or

$$\begin{aligned}\rho^T(-\tau) &= \rho(\tau), \quad \hat{A}^{(R)T}(-\tau) = \hat{A}(\tau), \\ \hat{H}^T(-\tau) &= \hat{H}(\tau), \quad \hat{U}^T(-\tau_1, -\tau_2) = \hat{U}(\tau_2, \tau_1)\end{aligned}\tag{A.8}$$

We note that the invariance of the functional integrals $|\psi\rangle, \{\psi|, \rho, \hat{U}, \hat{A}_{\delta}$ follows generally from the possibility to reverse the transformation (A.1) by a variable substitution. In contrast, the relations (A.3), (A.4) involve the invariance properties (A.2).

So far, all quantities have been real. We will now introduce a complex structure by considering two classes of functions of τ (including operators), namely those

which are either even or odd in τ . Even functions are considered as real, whereas the odd ones are purely imaginary. This relies on the isomorphism between a pair of real functions $(f_{\text{even}}, f_{\text{odd}})$ and the subclass of the complex functions $z = \text{Re } z + i \text{Im } z$, where $\text{Re } z$ is even and $\text{Im } z$ is odd, i. e. $(\text{Re } z, i \text{Im } z) \leftrightarrow (f_{\text{even}}, f_{\text{odd}})$. Complex conjugation is then equivalent to $(f_{\text{odd}} \rightarrow -f_{\text{odd}})$. Equivalently, the complex conjugation changes the sign of τ and we define its action as

$$\begin{aligned} |\psi(\tau)\}^* &= |\psi(-\tau)\} , \quad \{\psi(\tau)|^* = \{\psi(-\tau)|, \\ \rho(\tau)^* &= \rho(-\tau) , \quad \hat{H}(\tau)^* = \hat{H}(-\tau) \\ \hat{U}(\tau_2, \tau_1)^* &= \hat{U}(-\tau_2, -\tau_1) , \quad \hat{A}(\tau)^* = \hat{A}(-\tau) \end{aligned} \quad (\text{A.9})$$

Combining this definition with the action of the reflection θ (cf. eqs. (A.7), (A.8)) we recover well-known properties of quantum mechanics, namely

$$\begin{aligned} |\psi(\tau)\}^* &= \{\psi(\tau)|, \quad \rho^\dagger(\tau) = \rho(\tau), \quad \hat{H}^\dagger(\tau) = \hat{H}(\tau) \\ \hat{U}^\dagger(\tau_2, \tau_1) &= \hat{U}(\tau_1, \tau_2) = \hat{U}^{-1}(\tau_2, \tau_1) \end{aligned} \quad (\text{A.10})$$

With respect to this complex structure the reflection θ acts as hermitean conjugation. The euclidean time τ itself is odd and should therefore be considered as an imaginary quantity, $\tau = it$, t real. The self-consistency of this prescription is apparent by noting that for a wave function $|\psi(\varphi; \tau)\} = e^{-E\tau}\psi_0(\varphi)$ (E and ψ_0 real) the complex conjugate reads $\{\psi(\varphi; \tau)| = |\psi(\varphi; \tau)\}^* = (e^{-iEt})^*\psi_0(\varphi) = e^{iEt}\psi_0(\varphi) = e^{E\tau}\psi_0(\varphi) = |\psi(\varphi; -\tau)\}$. We note that $|\psi(0)\} = \{\psi(0)|$ and $\rho = \rho^T$ are real.

For an arbitrary operator $\hat{A}$ we define

$$\hat{A}^{(h)}(\tau) = \frac{1}{2} \left(\hat{A}(\tau) + \hat{A}^{(R)}(\tau) \right) , \quad \hat{A}^{(a)}(\tau) = \frac{1}{2} \left(\hat{A}(\tau) - \hat{A}^{(R)}(\tau) \right) \quad (\text{A.11})$$

From eqs. (A.8), (A.9) one finds $\hat{A}^{(R)}(\tau) = \hat{A}^\dagger(\tau)$ and we conclude that $\hat{A}^{(h)}$ is hermitean and $\hat{A}^{(a)}$ antihermitean

$$\hat{A}^{(h)\dagger}(\tau) = \hat{A}^{(h)}(\tau), \quad \hat{A}^{(a)\dagger}(\tau) = -\hat{A}^{(a)}(\tau) \quad (\text{A.12})$$

In compatibility with eq. (A.5) we conclude that all antihermitean operators must have purely imaginary expectation values. We emphasize that there is no a priori reason why antihermitean operators should not be associated with observables. In the sense of the original definition (2.5) their expectation values are simply odd with respect to the reflection of τ . The reflection symmetry alone does not enforce such expectation values to vanish. The situation is different for simultaneous τ -translation and reflection symmetry where in addition $\langle A(-\tau) \rangle = \langle A(\tau) \rangle$. In this case the expectation values of all odd observables or antihermitean operators vanish.

In conclusion, the discrete reflection symmetry introduces the complex structure characteristic for quantum mechanics: complex state vectors, hermitean density matrices and a hermitean Hamiltonian. The translation in τ is described by a unitary evolution operator.

B Analytic continuation

The definition of a complex conjugation as an involution in the space of τ -dependent functions does not yet specify the complex structure completely. In addition, we have to define the multiplication with complex numbers and the complex multiplication of functions. This is most easily done by constructing a mapping from the space of real functions of τ to the space of complex functions where the operations of complex conjugation and complex multiplication are implemented in the standard way. We will see that the analytic continuation of all functions $|\psi(\tau)\rangle$, $\hat{A}(\tau)$ etc. constitutes a map with all required properties once $\tau = it$ is considered as a pure imaginary variable.

We first show that analytic continuation of the functions of interest is indeed possible in the limit of continuous τ ($\epsilon \rightarrow 0$). With

$$\begin{aligned} (\partial/\partial\tau_2)^p \hat{U}(\tau_2, \tau_1) &= (-\hat{H}(\tau_2))^p \hat{U}(\tau_2, \tau_1), \\ (\partial/\partial\tau_1)^p \hat{U}(\tau_2, \tau_1) &= \hat{U}(\tau_2, \tau_1) (\hat{H}(\tau_1))^p \end{aligned} \quad (\text{B.1})$$

existing for all p , τ_2 and τ_1 the evolution operator $\hat{U}(\tau_2, \tau_1)$ is analytic both in τ_2 and τ_1 . In turn, $|\psi(\tau)\rangle$, $|\psi(\tau)|$ and $\hat{A}(\tau)$ are analytic functions of τ . As an example, we may represent $|\psi(\tau)\rangle$ as a Taylor series with real coefficients $a_n(\varphi)$, i.e. $|\psi(\varphi; \tau)\rangle = \sum_{n=0}^{\infty} a_n(\varphi) \tau^n$. Similarly, the symmetric and antisymmetric combinations read

$$\begin{aligned} |\psi_s(\tau)\rangle &= \frac{1}{2}(|\psi(\tau)\rangle + |\psi(-\tau)\rangle) \\ &= \sum_{m=0}^{\infty} a_{2m}(\varphi) \tau^{2m} = \sum_{m=0}^{\infty} (-1)^m a_{2m}(\varphi) t^{2m} \\ |\psi_a(\tau)\rangle &= \frac{1}{2}(|\psi(\tau)\rangle - |\psi(-\tau)\rangle) \\ &= \sum_{m=0}^{\infty} a_{2m+1}(\varphi) \tau^{2m+1} = i \sum_{m=0}^{\infty} (-1)^m a_{2m+1}(\varphi) t^{2m+1} \end{aligned} \quad (\text{B.2})$$

One sees that the complex conjugation (A.9) is compatible with the analytic continuation of the real function $|\psi(\tau)\rangle$ for $\tau \rightarrow it$. The descriptions in terms of the original real functions $|\psi(\tau)\rangle = |\psi_s(\tau)\rangle + |\psi_a(\tau)\rangle$ or the complex wave functions $|\psi(it)\rangle = |\psi_s(it)\rangle + |\psi_a(it)\rangle$ are completely equivalent – they are simply related to a change of variables. We repeat, however, that the use of the complex structure related to τ -reflection necessarily implies that τ must be purely imaginary. (Real values for τ are not compatible with this complex structure and should be used only in the language employed originally where all quantities are real. Obviously, the meaning of the word “real” depends on the complex structure used to distinguish between real and imaginary numbers. In absence of a complex structure all quantities are trivially “real”.)

The complex multiplication of two functions $|\psi_1(it)\rangle$ and $|\psi_2(it)\rangle$ is equivalent to the (real) multiplication of $|\psi_1(\tau)\rangle$ and $|\psi_2(\tau)\rangle$ in the original language

$$|\psi_3(\tau)\rangle = |\psi_1(\tau)\rangle|\psi_2(\tau)\rangle \leftrightarrow |\psi_3(it)\rangle = |\psi_1(it)\rangle|\psi_2(it)\rangle \quad (\text{B.3})$$

This is a direct consequence of the compatibility of analytic continuation with the product of two complex functions. The multiplication law (B.3) extends to matrix products $\hat{A}_1(it)\hat{A}_2(it)$ or $\hat{U}(it_2, it_1)|\psi(it_1)\rangle$. In the following we will always use the complex structure with the complex multiplication and adopt the notation

$$\begin{aligned} |\psi(t)\rangle &\equiv |\psi(it)\rangle, \quad \langle \psi(t)| \equiv \{\psi(it)| = |\psi(-it)\rangle = |\psi(t)\rangle^* \\ U(t_2, t_1) &\equiv \hat{U}(it_2, it_1), \quad A^{(M)}(t) \equiv \hat{A}(it) \end{aligned} \quad (\text{B.4})$$

In other words, $|\psi(t)\rangle$ is the analytic continuation of the functional integral $|\psi(\tau)\rangle$ given by eq. (2.9). In particular, for an exponentially decreasing $|\psi(\tau)\rangle = e^{-E\tau}\psi_0$ the state vector $|\psi(t)\rangle = e^{-iEt}\psi_0$ is an oscillatory complex function of t . We note that U is unitary

$$U^\dagger(t_2, t_1)U(t_2, t_1) = 1 \quad (\text{B.5})$$

and the time evolution conserves the norm of the complex state vector

$$\langle \psi(t)|\psi(t)\rangle = \langle \psi(\tau)|\psi(\tau)\rangle = 1 \quad (\text{B.6})$$

We associate the variable t with time and recover the Schrödinger equation (cf. (3.2))

$$i\partial_t|\psi\rangle = H|\psi\rangle \quad (\text{B.7})$$

At this point we should remark that our one-dimensional classical statistical system with next-neighbor interactions (1.1) has not a priori an interpretation as a description of a time evolution. In fact, one possibility is to view this system as a chain in space in equilibrium. In this case it describes infinitely many interacting degrees of freedom and there is no time evolution. The quantum mechanical discreteness is present, nevertheless, as well as the description in terms of operators and the possibility to introduce a complex structure. We have seen in appendix A that we can define operators and correlation functions in dependence on a complex parameter $\eta = e^{i\alpha t}$, e.g.

$$\hat{Q}_H(\eta) = e^{i\eta\hat{H}} \hat{Q} e^{-i\eta\hat{H}} \quad (\text{B.8})$$

Using the complex language (which is not familiar in this context) the relevant correlation functions for a chain in space correspond to purely imaginary η . Actually, the concepts of distance and geometry (in this case trivial) can be constructed from the η -dependence of the correlation function without assuming a space interpretation a priori [8].

A different interpretation of the probability distribution (1.1) concentrates on expectation values and correlations for real η . In this case η can be considered as a time parameter and our system describes the quantum-mechanical time evolution of

a single degree of freedom (not infinitely many as for the space interpretation). We emphasize that this interpretation is not characterized by a given deterministic time evolution equation. The concept of time arises from the “transport of information” between neighboring regions (“time regions” in this case) and is itself of a probabilistic nature! In this sense we may consider the dynamics of quantum mechanics as the result of an interpretation of time as a particular structure among observables in a classical statistical system.

Our definition of a complex structure within incomplete statistics allows the computation of expectation values of time-dependent observables $A(t)$ for a range of complex values of t . (The imaginary part of t should be within the local region $|\tau| \leq \bar{\tau}$ introduced in the formulation of our problem.) Although suggestive, it is not apparent in the present framework why our interpretation of the real world concentrates on real values of t . At the present stage it is not obvious why the values of the correlation functions at imaginary values of τ (real t) are much more important than their values of real τ . We believe that a satisfactory answer to these questions will shed more light on the basic origins of quantum mechanics and time.

If τ becomes a complex variable we may also consider analytic continuation directly in the functional integral. This can easily be achieved by choosing a purely imaginary ϵ in eqs. (2.2) and (2.4). Such a prescription brings us, however, to a path integral with phases for which the positivity of $\exp(-S)$ and therefore the classical probabilistic interpretation is lost. On the other hand, we could now release the restriction that the analytic continuation of $|\psi(t) \rangle$ leads to a positive real function $|\psi(\tau)\rangle$, opening the possibility to describe excited states. At the present stage we have not yet succeeded to overcome this conflict between the phases of quantum mechanics and the probabilistic interpretation of $\exp -S$. It seems worthwhile to investigate if there could exist more complex statistical systems for which the analytical continuation from euclidean to real time is consistent with a probabilistic interpretation.

References

- [1] A. Einstein, B. Podolski, N. Rosen, Phys. Rev. **47** (1935) 777
- [2] S. Weinberg, Phys. Rev. Lett. **62** (1989) 485
- [3] G. 't Hooft, gr-qc/9903084
- [4] J. S. Bell, Physical **1** (1964) 195
- [5] R. P. Feynman, Rev. Mod. Phys. **367** (1948) 20; Phys. Rev. **80** (1950) 440
- [6] J. Zinn-Justin, *Quantum Field Theory and Critical Phenomena*, (Oxford University Press 1996); S. Weinberg, *The Quantum Theory of Fields*, (Cambridge University Press 1995)

- [7] C. Wetterich, Nucl. Phys. **B314** (1989) 40
- [8] C. Wetterich, Nucl. Phys. **B397** (1993) 299
- [9] K. Osterwalder, R. Schrader, Comm. Math. Phys. **31** (1973) 83